

Connecting Rivers, Landscapes, People

Benwell Forest

Ecological Objectives & Hydrological Requirements

Justification Paper


Final: December 2014



NORTH CENTRAL


Catchment Management Authority

Connecting Rivers, Landscapes, People



Acknowledgement of Country

The North Central Catchment Management Authority acknowledges Aboriginal Traditional Owners within the region, their rich culture and spiritual connection to Country. We also recognise and acknowledge the contribution and interest of Aboriginal people and organisations in land and natural resource management.



Document control

Revision and distribution

Version no.	Description	Issued to	Issue date
1	Draft prepared by P. Beattie	North Central CMA SDL team	August 2014
2	Updated draft prepared by P. Beattie – includes feedback from stakeholders and CMA staff.	North Central CMA SDL team	September 2014
3	Final draft prepared by P. Beattie – includes minor amendments to reflect latest feedback from SDL team. E.g. discovery of colonial nesting in Benwell Swamp.	North Central CMA SDL team	21 October 2014
4	Final prepared by P. Beattie – includes expert review feedback from Terry Hillman and additional information to reflect updated project scope and latest investigations.	North Central CMA SDL team	11 November 2014
5	Final (version 2) prepared by P. Beattie. Includes minor amendments following business case submission.	North Central CMA SDL team	4 December 2014

Management

Organisation North Central Catchment Management Authority
 Author(s) Pam Beattie
 Number of pages 50
 Name of document Benwell Forest ecological objectives and hydrological requirements justification


This document should be referenced as North Central CMA (2014) *Benwell Forest ecological objectives and hydrological requirements*, Justification report prepared for the Sustainable Diversion Limit business case.

North Central Catchment Management Authority
 PO Box 18
 Huntly Vic 3551
 T: 03 5440 1800
 F: 03 5448 7148
 E: info@nccma.vic.gov.au
www.nccma.vic.gov.au

© North Central Catchment Management Authority, 2014


The North Central Catchment Management Authority wishes to acknowledge the Victorian and Commonwealth governments for providing funding for this publication through the National Action Plan for Salinity and Water Quality.

This is a working document, compiled from the best available information. It will be subject to revision in the future as new information becomes available. This publication may be of assistance to you, but the North Central Catchment Management Authority and its employees do not guarantee that the publication is without flaw of any kind, or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on information in this publication.



Contents

1. INTRODUCTION	6
1.1 Background	6
1.2 Workshop	6
2. Ecological values	8
2.1 Vegetation	8
2.2 Birds	11
2.3 Native fish	12
3. Condition	13
3.1 Prior to 2010-11	13
3.2 Post 2010-11	13
3.3 Ongoing expected flooding deficit	14
4. Ecological objectives and targets	16
5. HYDROLOGICAL REQUIREMENTS	22
5.1 Overview of requirements	22
5.2 Justification for hydrological requirements	24
6. PROPOSED WATER REGIME	26
7. BIBLIOGRAPHY	27
Appendix 1: Literature review of hydrological requirements	39




TABLES

Table 1.	Ecological Vegetation Classes (EVCs) present in Benwell Forest	9
Table 2.	Species of conservation significance recorded in the Guttrum and Benwell Forests	11
Table 3.	Bird species of conservation significance present in the Benwell Forest.....	11
Table 4.	Possible native fish species at Benwell Forest	12
Table 5.	Evaluation of mean daily flow series from Barham 1895-2009	14
Table 6.	Overarching ecological objectives and their justifications for Benwell Forest	17
Table 7.	Ecological objectives and targets for the Benwell Forest Environmental Works Project	19
Table 8.	Indicative hydrological requirements to achieve the Benwell Forest objectives	23
Table 9.	Hydrological requirements of wetland EVCs in Benwell Forest	24
Table 10.	Hydrological requirements of dominant River Red Gum EVCs in Benwell Forest.....	25
Table 11.	Waterbirds typical of Benwell Forest and their different breeding timeframes (North Central CMA 2009) .	48

FIGURES

Figure 1.	Benwell Forest water regime classes (Ecological Associates 2013)	8
Figure 2.	Ecological Vegetation Classes in Benwell Forest.....	10



1. INTRODUCTION

1.1 Background

The Commonwealth Department of Environment (DoE) is assessing a number of Environmental Works and Measures projects across the Murray Darling Basin that aim to achieve similar or better environmental outcomes using less water than previously estimated. The water savings generated by the projects could be used to increase the Sustainable Diversion Limit set under the Murray Darling Basin Plan by reducing the amount of water needed to be recovered from agricultural and urban use for the environment.

North Central Catchment Management Authority (North Central CMA) is currently developing business cases for two such projects – the Gunbower National Park Environmental Works Project and the Guttrum and Benwell Forests Environmental Works Projects.

This document summarises the **ecological values, objectives and targets of the Benwell Forest component of the Guttrum and Benwell Forests Environmental Works Project and the justification for the corresponding hydrological requirements** to achieve the objectives/targets.

The document has been divided into a series of chapters which documents the following:

- Values (i.e. what is there or what would we like there?) – this may include specific flagship species or broader ecological groupings depending on the ecological component being discussed e.g. Ecological Vegetation Classes, waterbird feeding guilds.
- Current condition and projected trajectories of condition.
- Ecological objectives/targets – these outline our goals for each ecological component considering their current condition and projected trajectories of condition.
- Hydrological requirements (i.e. what water regime is required to achieve the objectives/targets?) – based on scientific evidence where possible and/or practical experience.

The ecological components within this document have been based on those used for the Gunbower Forest National Park Environmental Works project and the broader Gunbower Forest TLM Icon Site – vegetation, birds and fish.

1.2 Workshop

A workshop was held on 31 July to obtain feedback on the ecological objectives and the required water regimes for the two projects mentioned above. Participants included expert ecologists, agency stakeholder representatives and North Central CMA staff including:

Kate Bennetts (Fire Flood and Fauna)


Rick Webster (Ecosurveys)

Marcus Cooling (Ecological Associates)

Clayton Sharpe (CPS Environmental)

Doug Frod (Pathways, Bushland and Environment)

Jack Smart (MDBA)



Emer Campbell (NCCMA)

Tim Hoogwerf (NCCMA)

Genevieve Smith (NCCMA)

Anna Chatfield (NCCMA)

Chris Corr (NCCMA)

Pam Beattie (NCCMA)

Andrea Keleher (DEPI)

Sharada Ramamurthy (DEPI)


Stephen Nicol (DEPI)

Paul Lacy (GMW)

Peter Foster (Parks Victoria)

Angela Gaynor (Parks Victoria)

This document captures the latest information relevant to the above aspects of the Benwell Forest Environmental Works Project, which considers the workshop discussions and incorporates the suggestions of participants following the workshop where appropriate.



2. ECOLOGICAL VALUES

2.1 Vegetation

Benwell Forest supports a diversity of vegetation, which can be described at a number of scales.

Landscape components

The Landscape Logic approach includes broad habitat types and how they are influenced by their position in the landscape. Within Benwell Forest, the habitat types include wetlands (semi-permanent) and River Red Gum forest (with flood dependent and flood tolerant understorey) (see diagram below) (Ecological Associates 2013).

Semi-permanent wetland habitat is provided in Benwell Forest through Benwell Swamp and Southwest Benwell Swamp (Ecological Associates 2013).





Figure 1. Benwell Forest water regime classes (Ecological Associates 2013)



Ecological Vegetation Classes

Ecological Vegetation Classes (EVCs) describe vegetation communities in Victoria using a combination of floristics, lifeform, position in the landscape and an inferred fidelity to particular environments. Each EVC includes a collection of floristic communities (i.e. groups based on co-occurring plant species) that occur across a bio-geographic range, and although differing in species, have similar habitat and ecological processes operating. Descriptions include canopy, understorey and groundcover species and each EVC is given a bioregional conservation status. Benchmarks (standard vegetation-quality reference points) are included and can be applied when carrying out vegetation quality assessments.

Across the Benwell Forest, 7 EVCs have been recorded. These are shown in the below table and illustrated in Figure 3. .

Table 1. Ecological Vegetation Classes (EVCs) present in Benwell Forest

Ecological Vegetation Class	Bioregional Conservation Status	Extent (ha)
Floodway Pond Herbland / Riverine Swamp Forest Complex (EVC 945)	Depleted	41
Grassy Riverine Forest (EVC 106)	Depleted	119
Riverine Grassy Woodland (EVC 295)	Vulnerable	9
Riverine Swamp Forest (EVC 814)	Depleted	339
Sedgy Riverine Forest (EVC 816)	Depleted	51
Spike-sedge Wetland (EVC 819)	Vulnerable	13
Floodplain Riparian Woodland (EVC 56)	Depleted	10

Source: Ecological Associates (2013).





Figure 2. Ecological Vegetation Classes in Benwell Forest

Species

More than 75 species of native flora have been recorded in Guttrum and Benwell State Forests in a single survey, including 20 rare or threatened species of which 19 were listed under the FFG Act (1988) (Biosis 2014). It is expected that many more species will be recorded in the forest during future investigations, similar to those species identified in the nearby Gunbower Forest (Bennetts 2014).

Flora species of conservation significance recorded in the Guttrum and Benwell Forests are shown below.




Table 2. Species of conservation significance recorded in the Guttrum and Benwell Forests

Flora species	Status		
	FFG	EPBC	VROTS
Branching groundsel (<i>Senecio cunninghamii</i> var. <i>cunninghamii</i>)			Rare
Flat-top saltbush (<i>Atriplex lindleyi</i> subsp. <i>lindleyi</i>)			Poorly known
Mealy saltbush (<i>Atriplex pseudocamanulata</i>)			Rare
Native peppergrass (<i>Lepidium pseudohyssopifolium</i>)			Poorly known
Smooth elachanth (<i>Elachanthus glaber</i>)			Rare
Smooth minuria (<i>Minuria integerrima</i>)			Rare
Stiff groundsel (<i>Senecio behrianus</i>)	Listed as threatened	Endangered	Endangered

Source: North Central CMA (2014a)

FFG – Flora and Fauna Guarantee Act 1988, EPBC – Commonwealth Environment Protection and Biodiversity Conservation Act 1999, VROTS –Victorian Rare or Threatened Species Advisory Lists

2.2 Birds


The Benwell Forest provides habitat for native birds including terrestrial and waterbird species. Benwell Forest provides habitat for the conservation significant Grey-crowned babbler which nests in River Red Gum at the southern edge of Benwell Forest. Several bird species of conservation significance occur across Benwell Forest and are summarised here (from Atlas of Victoria Wildlife).

Table 3. Bird species of conservation significance present in the Benwell Forest

Bird species	Status		
	FFG	EPBC	VROTS
Australasian bittern (<i>Botaurus poiciloptilus</i>)	Listed as threatened	Endangered	Endangered
Azure kingfisher (<i>Alcedo azurea</i>)			Near threatened
Brown treecreeper – south eastern subspecies (<i>Climacteris picumnus victoriae</i>)			Near threatened
Eastern great egret (<i>Ardea modesta</i>)			Vulnerable
Grey-crowned babbler (<i>Pomatostomus temporalis</i>)	Listed as threatened		Endangered
Intermediate egret (<i>Ardea intermedia</i>)	Listed as threatened		Endangered
Little egret (<i>Egretta garzetta</i>)	Listed as threatened		Endangered
Nankeen night heron (<i>Nycticorax caledonicus</i>)			Near threatened
Pied cormorant (<i>Phalacrocorax varius</i>)			Near threatened
Royal spoonbill (<i>Platalea regia</i>)			Near threatened
White-bellied sea eagle (<i>Haliaeetus leucogaster</i>)	Listed as threatened		Vulnerable

Source: North Central CMA (2014)

FFG – Flora and Fauna Guarantee Act 1988, EPBC – Commonwealth Environment Protection and Biodiversity Conservation Act 1999, VROTS –Victorian Rare or Threatened Species Advisory Lists



Colonial waterbird nests (thought to have been constructed in response to the 2010/11 floods) have recently been observed in Southwest Benwell Swamp (G. Smith, pers. comm, 16 October 2014). Based on the size of the nests, these may have been constructed by cormorants.


2.3 Native fish

Information on native fish within or using the Benwell Forest floodplain and wetlands is limited. Three species of conservation significance are expected to use the area – Murray cod, Golden perch and Silver perch (Ecological Associates 2013).

While fish survey data is unavailable, it is possible that a similar fish community exists in these forests to that which has been recorded for Gunbower Forest (see below). This is due to the similar habitat availability (wetlands within River Red Gum forest floodplain), adjacent waterway (River Murray) and the relatively close proximity of these forest systems to each other. Confirmation of the validity of this assumption is required by fish ecologists.

Table 4. Possible native fish species at Benwell Forest (based on those known and expected to occur at Gunbower Forest including the River Murray)

Group	Species Name	Common Name	EPBC	FFG
KNOWN SPECIES at Gunbower Forest				
Long-lived apex predators	<i>Maccullochella peelii peelii</i>	Murray cod	Vulnerable	Vulnerable
	<i>Maccullochella macquariensis</i>	Trout cod	Endangered	Critically Endangered
Flow dependent specialists	<i>Macquaria ambigua</i>	Golden perch		Near threatened
	<i>Bidyanus bidyanus</i>	Silver perch	Critically Endangered	Vulnerable
Foraging generalists	<i>Retropinna semoni</i>	Australian smelt		
	<i>Hypseleotris spp.</i>	Carp gudgeon (group)		
	<i>Melanotaenia fluviatilis</i>	Murray-Darling rainbowfish		Vulnerable
	<i>Philypnodon grandiceps</i>	Flathead gudgeon		
	<i>Philypnodon macrostomus</i>	Dwarf flat-headed gudgeon		
	<i>Tandanus tandanus</i>	Freshwater catfish		Endangered
	<i>Nematalosa erebi</i>	Bony bream		
	<i>Craterocephalus stercusmuscarum fulvus</i>	Un-specked hardyhead		
	<i>Craterocephalus stercusmuscarum</i>	Fly-specked hardyhead		
EXPECTED SPECIES at Gunbower Forest				
Foraging generalist	<i>Macquaria australasica</i>	Macquarie Perch	Endangered	Endangered
Floodplain specialists	<i>Nanoperca australis</i>	Southern Pygmy Perch		
	<i>Mogurnda adspersa</i>	Southern Purple-spotted Gudgeon		Regionally Extinct
	<i>Galaxias rostratus</i>	Flat-headed Galaxias		Vulnerable
	<i>Ambassis agassizii</i>	Olive Perchlet		Regionally Extinct
	<i>Craterocephalus fluviatilis</i>	Murray Hardyhead	Endangered	Critically Endangered



3. CONDITION

3.1 Prior to 2010-11

For ten years prior to 2010-11, the dry climatic conditions resulted in below average inflows and therefore lower water availability. Prior to the natural flood event in 2010-12, the Benwell Forest experienced inadequate, flooding compared to natural (with some unknown volumes likely to have been supplied through natural flooding in 2000 and 2003-2005 (K. Bennetts, pers. comm. August 2014). It was not possible to deliver environmental water to these sites due to a lack of appropriate infrastructure. Encroachment of terrestrial vegetation into areas formerly occupied by flood dependent species occurred during this period. While field data is limited, it is assumed that similar to other forests with inadequate flooding, the condition of the Benwell Forest declined during the last drought period.

In 2007, investigations noted that Benwell Forest had changed to a much younger and denser forest due to the altered frequency, duration and depth of flooding (SKM 2007).

The dry climatic conditions for ten years prior to 2010-11 are expected to have reduced floodplain productivity and access to food and habitat for native fauna. This potentially led to a decline in fauna populations and their resilience to additional stressors (Horrocks et al unpubl.). This was most evident for colonial waterbird populations — the extended periods between large floods that support large-scale breeding opportunities posed a key threat to the viability of existing populations (Murray Darling Basin Authority 2012).


Ecological processes required to sustain native fish populations, such as connectivity to the floodplain for breeding and recruitment, is also likely to have been hindered (Ecological Associates 2010).

3.2 Post 2010-11

In 2010-11, the Benwell Forest received extensive inundation from high River Murray flows. It is expected that the condition of the vegetation community improved as a result of the natural flooding.

The latest vegetation assessment (Bennetts 2014) found Benwell Forest supports considerable areas of floodplain forest with relatively intact understorey, rare and threatened species and stands of large old trees. However the structure of the forest ecosystem has been altered and degraded in some parts through a number of disturbances – river regulation, timber harvesting, cattle grazing, weed invasion and quarrying. The current condition is summarised below (Bennetts 2014):

- Wetlands are impacted by cattle grazing, with some wetland vegetation trampled or physically uprooted and pugging evident. The more palatable wetland flora are absent e.g. River Swamp Wallaby Grass. River Red Gum saplings have encroached on the wetland areas and resulted in less open water habitat. Medic (*Medicago* spp.) is one of the most abundance weed species, forming mono-specific patches in the wetlands and reducing the biodiversity in these locations. Perennial emergent macrophytes are limited to stands of *Phragmites australis* in the understorey of the River Red Gum forest and are absent elsewhere (Ecological Associates 2013).
- Large areas of River Red Gum forest are dominated by small (i.e. < 50 cm diameter at breast height (DBH)) trees due to long-term forestry practices – progressive removal of the tallest and straightest trees, ringbarking



the irregular trees and thinning regrowth. Most of the mature trees show signs of drought stress in their canopies (e.g. epicormic growth) and their ongoing health may be impacted by competition from the surrounding dense stands of regrowth. Some mature trees have recently died – potentially in response to the Millennium drought post long-term river regulation. In the southeast corner of Benwell Forest, some River Red Gums have died – presumably from drowning following extended ponding of water against the boundary levee.

- Between the wetlands and elevated River Red Gum woodland, are large forest areas with relatively intact understories that support characteristic, delicate and rare and threatened species.
- The old sand quarries are significantly disturbed and even though they currently retain water, they have limited ecological value.

A more detailed condition assessment is being undertaken across the forest and results will soon be available.

The flooding response from waterbirds after 2010-12 is unknown, however is expected to have been limited, primarily due to the loss of wetland habitat within the forests.

3.3 Ongoing expected flooding deficit

Similar to the nearby Gunbower Forest, the Benwell Forest is located in an area of low rainfall and high evapotranspiration. The average annual rainfall is less than 400 millimetres per year with evapotranspiration of around 1,700 mm/y. This creates a significant water deficit and stressor to the forest, particularly in years where there are no forest inflows to maintain health of vegetation communities (Murray Darling Basin Authority 2012).


The hydrology of these forests has changed substantially because of the regulation and diversion of River Murray flows, resulting in a reduction in the frequency and duration of flooding (see below). Recent modelling (Ecological Associates 2013) demonstrates that, under current river operations, intervention is needed to maintain functioning floodplain ecosystems within the Benwell Forest (Ecological Associates 2013).

Table 5. Evaluation of mean daily flow series from Barham 1895-2009


Flow threshold exceeded (ML/d)	Natural conditions		Benchmark conditions (including The Living Murray works)		Basin Plan (2750GL)	
	Median frequency (events/100yrs)	Median duration (days)	Median frequency (events/100yrs)	Median duration (days)	Median frequency (events/100yrs)	Median duration (days)
>15GL/day	100	175	80	75	91	115
>25GL/day	82	100	45	75	60	85
>33GL/day	25	40	8	49	9	49

Source: Ecological Associates 2013 (actual figures may vary slightly due to graph interpretation)

According to Ecological Associates (2013) and Gippel (2014), flows that inundate the semi-permanent wetlands now occur in 61% of years rather than the 90% of years that would have occurred naturally and last for 1 to 4 months (interquartile range at 21,000 ML/d) instead of 3 to 6 months. The reduction in flood duration and frequency and the longer periods between events has made wetland habitat less persistent and reliable. Red gum has encroached on the wetland areas in the Benwell Forest (Ecological Associates 2013).



Flows that inundate the River Red Gum forest with flood dependent understorey now occur in 37% of years rather than the 70% of years that is modelled under natural conditions (Gippel 2014). Typical events last for 1 to 3 months instead of 2 to 5 months (interquartile range at 27,000 ML/d). The reduced flooding frequency and duration has reduced plant productivity so that wetland, canopy and understorey vegetation has less frequent and shorter growing periods and provides less food resources and physical habitat for fauna. The brief and less frequent flooding has reduced habitat opportunities, breeding events for wetland fauna including waterbirds, foraging and nursery habitat opportunities for riverine fish and contributions of carbon to the river channel food web (Ecological Associates 2013). This decline in overall condition is expected to continue under future regulated river conditions.



4. ECOLOGICAL OBJECTIVES AND TARGETS

Ecological objectives and targets describe the intended outcomes of environmental water delivery. They contribute towards achieving the higher level goals (vision and management goal) including:

Benwell Forest vision:

To maintain and improve the Benwell Forest by restoring a more natural water regime that supports a variety of aquatic habitats and a range of native flora and fauna species.

Management goal for Benwell Forest Flood Enhancement Project:

Reinstate a more natural water regime that protects and enhances the ecological values within the Benwell Forest.

Two levels of objectives and targets have been developed (Table 3). The overarching objectives capture the primary, higher level aims of the project, while the detailed objectives break the overarching objectives down into the various ecological components i.e. they describe what a 'healthy' community may include, which can then be linked to monitoring methods and reporting against targets.

The ecological objectives and targets for the Benwell Forest project are based on the key values of the site (in line with the Victorian Waterway Management Strategy) including its:

- mosaic of habitat (e.g. foraging areas) required for macro/micro invertebrates, frogs, fish and waterbirds across multiple feeding guilds.
- ability to support native frog, native fish and waterbird breeding.
- ability to support species listed under international agreements (JAMBA, CAMBA), the *Environmental Protection Biodiversity Conservation Act* (1999) and the *Flora and Fauna Guarantee Act* (1988).
- ability to encourage recruitment of channel specialist native fish such as Murray Cod through floodwater inputs to the River Murray (Mallen-Cooper et al. 2014).

The objectives also take into consideration the:

- Hydrological changes in the system – the current and projected (under the Basin Plan) deficit in the water regime compared to natural inundation. Refer to previous section 'Ongoing expected flooding deficit'.
- Current condition and trajectory of each value and therefore whether intervention is required. Refer to previous section.
- Inter-dependencies within the Benwell Forest site and between local forest floodplains (e.g. Guttrum Forest, Campbells Island). For example, the success of colonial nesting waterbird breeding in Benwell Forest's Southwest Benwell Swamp is dependent in part on the availability and security of foraging areas. Therefore objectives around colonial waterbird breeding in Benwell Forest is inter-linked with the objectives around inundating River Red Gum to provide foraging areas for waterbirds in the Benwell Forest.

The overarching ecological objectives for the Benwell Forest project and the justification for each are shown below.





Table 6. Overarching ecological objectives and their justifications for Benwell Forest

Overarching objective	Justification
<p>Improve the health of semi-permanent wetlands</p>	<ul style="list-style-type: none"> ▪ Provides a mosaic of habitat (e.g. marshland, herbland, open water and reed bed plant communities) for aquatic and amphibious plants to reproduce, which provide food for waterbirds (Ecological Associates 2013). ▪ Provides food and habitat for micro/macro invertebrates, frogs and small fish (which provide food resources for waterbirds e.g. large wading birds and piscivores). ▪ Receding flood water in summer provides habitat for migratory wading birds that pick over invertebrates in drying mud (Ecological Associates 2013). ▪ Maintains the health and distribution of the existing River Red Gum population, which provide nesting, feeding and breeding habitat for fauna, as well as organic inputs to the wetland (North Central CMA 2012). ▪ Ensures viable seed bank and invertebrate egg bank is maintained (North Central CMA 2012). ▪ Wet/dry cycle consolidates soils, allows sedimentation processes to occur, mineralises organic matter, supports microbial and planktonic productivity and overall biodiversity in the long-term (North Central CMA 2012; Ecological Associates 2013). ▪ Wetland vegetation provides shelter, nesting habitat and nesting materials for breeding waterbirds. E.g. dense macrophyte beds are important for cryptic waterbirds like Australasian bittern, purple swamp hen and black-tailed native hen. Dense reed beds provide nesting habitat for swamp harrier and growling grass frog. Marshy areas with semi-emergent vegetation are important to grebes and dabbling ducks e.g. Great crested grebe and Australasian grebe. Fringing red gum forest provides habitat for colonial nesting waterbirds to build platform nests (Ecological Associates 2013).
<p>Healthy wetland bird community across Benwell Forest through improved access to food and habitat that promotes breeding and recruitment</p>	<ul style="list-style-type: none"> ▪ Rehabilitating a diversity of foraging habitat supports a high carrying capacity of waterbirds across Benwell Forest including those residing/breeding in the forest and those residing/breeding in nearby areas, which use these forests as part of their broader foraging areas (e.g. Colonial nesting species). This will promote a diverse waterbird community from a range of feeding guilds. ▪ Waterbird breeding success is correlated with the foraging area available. For example, breeding waterbirds within Gunbower Forest have been reported to move on a daily basis to the adjacent Koondrook Perricoota area for foraging (North Central CMA 2009). ▪ The wetlands are expected to support a variety of waterbird breeding events. ▪ Provision of appropriate habitat will support recruitment of waterbirds at a landscape scale.
<p>Healthy River Red Gum communities</p>	<ul style="list-style-type: none"> ▪ Provides food and habitat for micro/macro invertebrates (which are food resources for frogs, fish and waterbirds). ▪ The recently flooded understorey provides seeds, fruit and forage for granivores such as finches, cockatoos, galah, lorikeet and budgerigar, the frugivorous emu and herbivorous swamp wallaby (Ecological Associates 2013). ▪ Flood dependent understorey potentially reduced in extent prior to 2010, due to terrestrial encroachment.



Overarching objective	Justification
	<ul style="list-style-type: none"> ▪ The trees directly support nectivorous and omnivorous birds such as honeyeaters and wattlebird (Ecological Associates 2013). ▪ Helps maintain wetland productivity by providing organic inputs to the water column (e.g. carbon and nutrient inputs from leaf litter). ▪ Provides nesting material ((including hollows) and roosting habitat for waterbirds. ▪ Tree canopy was observed as still recovering from the drought in some locations during recent field observations (K. Bennetts pers. com July 2014).
<p>Promote recruitment of the local River Murray channel specialist native fish community^ by increasing opportunities to access productive floodplain outflows from Benwell Forest.</p>	<ul style="list-style-type: none"> ▪ Provides organic inputs from the floodplain to the waterways, promoting instream productivity (North Central CMA 2014a). ▪ Supports recruitment of channel specialist fish through improved instream productivity i.e. higher survival of larvae as floodwaters (high in phytoplankton and zooplankton) recede back into channels (Mallen-Cooper et al. 2014). ▪ Recruitment occurs when floodplains are inundated, which increases productivity and larval survival. Likely applies to all fish species to some degree (Mallen-Cooper et al. 2014). ▪ Improved access to food resources from the floodplain will enhance the health and potentially the resilience of channel specialist fish species.

^ Mainly large bodied native fish that spawn within channel habitats (such as Murray Cod, Golden Perch and Silver Perch).

The full range of ecological objectives and targets for the project are shown below.






Table 7. Ecological objectives and targets for the Benwell Forest Environmental Works Project


Objectives (by 2040)	Targets (by 2040)	Applicable values
SEMI-PERMANENT WETLANDS		
<ul style="list-style-type: none"> ▪ Overarching: Improve the health of semi-permanent wetlands 	<ul style="list-style-type: none"> ▪ S1 (Overarching): >95% of semi-permanent wetlands with a water regime that maximises healthy condition. ▪ S2: At least 75% of wetland transects with ‘moderate to excellent’ vegetation condition as defined by the TLM wetland and floodplain condition assessment categories. 	<ul style="list-style-type: none"> ▪ Rare freshwater meadow with diverse habitats ▪ Nationally vulnerable Growling Grass Frog. ▪ Waterbird feeding and breeding habitat.
<ul style="list-style-type: none"> ▪ Achieve an appropriate cover and diversity of species characteristic of the Plant Functional Groups found in the semi-permanent wetlands. 	<ul style="list-style-type: none"> ▪ S3: Plant Functional Groups 1-7 have >50% of total cover occupied by at least 2/3 of all species possible within these Plant Functional Groups. (See Monitoring & Evaluation Plan for further details). 	
<ul style="list-style-type: none"> ▪ Reduce River Red Gum encroachment in semi-permanent wetland areas. 	<ul style="list-style-type: none"> ▪ S4: River Red Gum encroachment is absent. 	
<ul style="list-style-type: none"> ▪ Provide suitable habitat for the threatened (EPBC listed) Growling grass frog. 	<ul style="list-style-type: none"> ▪ S5: Presence of habitat suitable for the EPBC listed Growling grass frog. i.e. water with diverse habitat including emergent, submerged and floating vegetation within the September to March breeding period in at least 7 years in 10 e.g. Reed Bed Swamp, Southwest Benwell Swamp. (See Monitoring & Evaluation Plan for further details). 	
<ul style="list-style-type: none"> ▪ Maintain and where possible increase the current diversity of threatened flora species. 	<ul style="list-style-type: none"> ▪ S6: >50% of threatened species previously recorded observed. (See Monitoring & Evaluation Plan for further details). 	
<ul style="list-style-type: none"> ▪ Reduce the area of high threat weed species. 	<ul style="list-style-type: none"> ▪ S7: High threat exotic plants absent in >90% of total cover. 	
NATIVE BIRDS		
<ul style="list-style-type: none"> ▪ Overarching: Healthy wetland bird community across Benwell Forest through improved access to food and habitat that promotes breeding and recruitment 	<ul style="list-style-type: none"> ▪ B1 (Overarching): Successful colonial waterbird breeding in at least 3 years in 10 (for a range of species – egrets, cormorants, herons). ▪ B2: Successful waterfowl breeding in at least 7 years in 10. 	<ul style="list-style-type: none"> ▪ Bird species of conservation significance. ▪ Waterbird feeding and breeding habitat.



Objectives (by 2040)	Targets (by 2040)	Applicable values
<ul style="list-style-type: none"> Support a suite of waterbirds including waterfowl, colonial waterbirds and other wetland dependent species. 	<ul style="list-style-type: none"> B3: >50% of all waterbird species expected to occur observed over a ten-year period. 	
<ul style="list-style-type: none"> Provide foraging areas for colonial nesting waterbirds in Benwell Forest and potentially elsewhere (e.g. lower Gunbower Forest). 	<ul style="list-style-type: none"> B4: >60% of the floodplain inundated for colonial waterbird foraging in 8 years in 10. 	
<ul style="list-style-type: none"> Maintain and where possible increase the current diversity of threatened wetland bird species. 	<ul style="list-style-type: none"> B5: Presence of water in swamps with emergent vegetation e.g. Reed Bed Swamp, such as <i>Typha spp.</i> (bullrush), <i>Phragmites spp.</i> (reeds) and <i>Eleocharis spp.</i> (sedges) in at least 7 years out of 10. 	
RIVER RED GUM FOREST		
<ul style="list-style-type: none"> Overarching: Healthy River Red Gum communities. 	<ul style="list-style-type: none"> R1 (Overarching): >80% of River Red Gum forest with flood dependent understorey with a water regime that maximises healthy condition. R2: Range of age classes exist for River Red Gums in at least 75% of surveyed areas. 	<ul style="list-style-type: none"> FFG Act listed River Red Gum Grassy Woodland ecological community. 23 threatened flora species
<ul style="list-style-type: none"> Achieve an appropriate cover and diversity of species characteristic of the Plant Functional Groups found in the River Red Gum forest understorey. 	<ul style="list-style-type: none"> R3: River Red Gum with flood dependent understorey – <ul style="list-style-type: none"> Plant Functional Groups 2-7 have >50% of total cover occupied by at least 2/3 of all species possible within these Plant Functional Groups. (See Monitoring & Evaluation Plan for further details). 	
<ul style="list-style-type: none"> Maximise the proportion of trees with healthy canopy condition in the River Red Gum forests. 	<ul style="list-style-type: none"> R4: >75% of surveyed trees with ‘healthy’ canopy condition - crown condition index score of 4 or greater. 	
<ul style="list-style-type: none"> Maintain and where possible increase the current diversity of threatened flora species. 	<ul style="list-style-type: none"> R5: >50% of threatened flora species previously recorded observed. 	
<ul style="list-style-type: none"> Reduce the area of high threat weed species. 	<ul style="list-style-type: none"> R6: High threat exotic plants absent in >90% of total cover. 	
NATIVE FISH		



Objectives (by 2040)	Targets (by 2040)	Applicable values
<ul style="list-style-type: none"> ▪ Overarching: Promote recruitment of the local River Murray channel specialist native fish community by increasing opportunities to access productive floodplain outflows from Benwell Forest. 	<ul style="list-style-type: none"> ▪ F1: Commonly occurring large-bodied, channel specialist native fish species (Murray Cod and Golden Perch) occur every year in local River Murray surveys and include a range of age and size classes. 	<ul style="list-style-type: none"> ▪ Diverse fish community similar to that recorded around Gunbower Forest expected. ▪ EPBC listed Murray Cod in the River Murray.



5. HYDROLOGICAL REQUIREMENTS

5.1 Overview of requirements

The indicative hydrological requirements for each ecological component described through the objectives are shown below. The justification for these water requirements is provided below and is based on a substantial literature review as well as input by expert ecologists.

These hydrological requirements have been used to inform the proposed water regime for the project outlined in the next chapter.





Table 8. Indicative hydrological requirements to achieve the Benwell Forest Environmental Works Project objectives

Ecological objective		Hydrological Objectives										
		Recommended number of events in 10 years			Tolerable interval between events once wetland is dry (months)			Duration of ponding (months)			Preferred timing of inflows	Depth (m)
		Min	Opt	Max	Min	Opt	Max	Min	Opt	Max		
Improve the health of semi-permanent wetlands)		6	9	10	1	6	36	3	6	8	Winter/ spring	Often <50cm. The general depth of Benwell Swamp is 0.5m and 0.7m in Southwest Benwell Swamp.
Healthy bird community across Benwell Forest through improved access to food and habitat that promotes breeding and recruitment	Egrets	3	4	5	12	18	24	10*	12	12	Late winter/spring/early summer	Not critical but maintain depth during breeding and have gradual changes
	General waterbirds (not colonial nesting species)	3	5	10	12	18	24	4	6	12	Late winter/ spring/ early summer	Maximise area up to 30cm deep. Need to fluctuate depth over time to promote wetland productivity.
Healthy River Red Gum communities		6	7	8	-	-	36	4	4	7	Winter/ spring	Not critical for adult River Red Gums. Varies for understorey. Some understorey sp. prefer shallow depths <10cm during active growth but can tolerate deeper immersion for short periods.
Promote recruitment of the local River Murray channel specialist native fish community by increasing opportunities to access productive floodplain outflows from Benwell Forest.		6	7	8	-	-	36	4	4	7	Outflows to River Murray in spring/summer after temperature and flow cued spawning occurs in the channel.	Sharp drop in water level required to provide a fish exit cue in late spring/summer for any fish that have entered the floodplain. Hypothesis: 0.3m over 48 hrs. This will also promote organic matter export.

* Duration of inundation is 5-7 months for cormorants (North Central CMA 2009).



5.2 Justification for hydrological requirements

Evidence to support the hydrological requirements for each ecological objective is outlined below. This includes a combination of primary and secondary literature/reports, as well as input by expert ecologists. A literature review is available in the Appendix.

Semi-permanent wetlands (Overarching objective: Improve the health of semi-permanent wetlands)

According to recent surveys (Bennetts 2014; Biosis & Bennetts 2014), semi-permanent wetlands in Benwell Forest primarily consist of Floodway Pond Herbland/Riverine Swamp Forest Complex (EVC 945) with smaller areas of Spike-sedge Wetland (EVC 819). The water regime for this objective therefore reflects that required by the dominant wetland EVC, which is similar to that required by the less extensive wetland EVC as shown below (Fitzsimons et al. 2011).

See Appendix 1 for further details on wetland water requirements.

Table 9. Hydrological requirements of wetland EVCs in Benwell Forest

Ecological Vegetation Class	Natural flood frequency	Critical interval between events (years)	Minimum duration (months)
Floodway Pond Herbland/Riverine Swamp Forest (EVC 945)	6-10 in 10	3	3-8
Spike-sedge wetland (EVC 819)	5-10 in 10	4	1-4


Source: Fitzsimons et al. 2011

Waterbirds (Overarching objective: Healthy wetland bird community across Benwell Forest through improved access to food and habitat that promotes breeding and recruitment)

This objective includes two sets of hydrological requirements – those needed for Egrets (an indicator species used to represent the water requirements of the broader group of colonial nesting waterbirds) and those needed for non-colonial waterbirds.

Achieving successful colonial waterbird breeding events requires a specific water regime in addition to that required by the general waterbird community. Egrets (Great Egret (*Ardea alba*), Intermediate Egrets (*Ardea intermedia*) and Little Egrets (*Egretta garzetta*)) are known to be sensitive to flooding conditions when breeding. For example, they require the longest flood duration of any waterbird in these forests. It is assumed that implementing an operating strategy that leads to successful breeding of egrets, will also enable other waterbird species to breed successfully. This is based on information obtained through a waterbird breeding requirement workshop, which found providing conditions suitable for colonial species will also provide suitable breeding conditions for a wide range of other species such as ducks, grebes, swampheens and herons (North Central CMA 2009).

An expert bird ecologist has confirmed the hydrological requirements outlined above, during the Ecological Objectives refinement workshop (R. Webster, pers. comm. July 2014). Appendix 1 includes details on the water regime required for general waterbirds.



River Red Gum communities (Overarching objective: Healthy River Red Gum communities)

The River Red Gum forest with flood dependent understorey across both sites is predominantly Riverine Swamp Forest (EVC 814). In Benwell Forest, a substantial portion of the eastern floodplain is Grassy Riverine Forest (106) (Bennetts 2014), with Riverine Swamp Forest dominating the western portion. The water regime has therefore been based on the Riverine Swamp Forest EVC. The optimum duration of ponding has been chosen to reflect that suitable for both the Riverine Swamp Forest and Grassy Riverine Forest EVCs as shown below.

Table 10. Hydrological requirements of dominant River Red Gum EVCs in Benwell Forest

Ecological Vegetation Class	Natural flood frequency	Critical interval between events (years)	Minimum duration (months)
Riverine Swamp Forest (EVC 814)	6-8 in 10	3	4-7
Grassy Riverine Forest (EVC 106)	5-10 in 10	4	1-4

Source: Fitzsimons et al. 2011


Note: Given the forest has experienced stress from the drought and is still recovering, there will be a gradual build up to this water regime as recommended by an expert vegetation ecologist (Bennetts, K. pers. comm. August 2014).

Native fish (Overarching objective: Promote recruitment of the local River Murray channel specialist native fish community by increasing opportunities to access productive floodplain outflows from Benwell Forest.)

Native channel fish access to productive floodplain outflows (containing organic matter) from the Benwell Forest floodplain is provided through broad inundation of the River Red Gum forests. The hydrological requirements for this objective therefore reflect that defined for River Red Gum forest with flood dependent understorey (Riverine Swamp Forest EVC 814 and Grassy Riverine Forest EVC 106). See Appendix 1 for further details on native fish requirements.

In general, the only channel species that require a rise in flow or a flood to spawn are the flow-cued spawners, golden perch and silver perch. All other species reliably spawn each year in response to rising temperatures in spring and summer (Mallen-Cooper et al. 2013).

This means that to maximise the contribution of floodplain outflows to in-channel recruitment for species such as golden perch, the outflows should occur, where possible, after variable channel flows that trigger spawning. For other species (e.g. Murray Cod), the floodplain outflows should occur in spring/summer after migration and temperature cued spawning has occurred.



6. PROPOSED WATER REGIME

The water regime for the Benwell Forest has been based on consideration of:

- The relationship between objectives for example, healthy wetland birds are partly provided through inundation of the semi-permanent wetland habitat. Access of channel fish to floodplain waters depends on the inundation of broad areas of River Red Gum floodplain that can then drain into the River Murray.
- Risk management – in terms of broader floodplain watering, the water regime is conservative to manage any risks associated with re-introducing regular watering to a forest that has experienced a flooding deficit for an extended period.

Therefore the water regime for the Benwell Forest may be defined for two scenarios as below.


Scenario 1 –River Red Gum floodplain:

- Frequency: 8 years in 10
- Duration of inundation: 4 months
- Timing: Winter/spring

Scenario 2 – Semi-permanent wetland watering:

- Frequency: 9 years in 10
- Duration of inundation: 6 months
- Timing: Winter/spring, with drying (drawdown of water level) in late summer/autumn.
- Depth: fluctuate over time, inundate to Full Supply Level in some years.

Note: if a substantial bird breeding event occurs (determined on an event by event basis), then ongoing environmental water deliveries (top-ups) may be required and will be provided as necessary. Detailed monitoring and advice from specialist ecologists will be used to determine the water requirements of each colony.



7. BIBLIOGRAPHY

Akeroyd, M. D., Tyerman, S. D., Walker, G. R. & Jolly, J. D. 1998. 'Impact of flooding on the water use of semi-arid eucalypts', *Journal of Hydrology*, vol. 54, pp. 104-117.

Allen, G., Midgley, S. and Allen, M. 2003, *Field Guide to Freshwater Fishes of Australia*, 2nd ed., CSIRO Publishing, Melbourne.

Amphibian Research Centre. 2014, *Frogs of Australia*, www.frogs.org.au, accessed May 2014.

Anderson, B. G., Gippel, C. J., Cooling, M. P., Lloyd, L. N. and Kerr, G. 2007, *Gunbower Creek Environmental Flows Study, Issues Paper*, report by Water Technology in association with Fluvial Systems, Lloyd Environmental and Ecological Associates for North Central Catchment Management Authority, Huntly.

Atkins, B. P., Lloyd, L. N. and Nikolaou, N. 1991, *The Hydrological Characteristics of Gunbower Forest*, Draft Report of the Integrated Watering Strategy (ed. DfCa Environment).

Atlas of Living Australia. 2014, *Web Pages on Various Frog Species*, accessed May 2014, <http://www.ala.org.au>

Australian Ecosystems. 2009, *Gunbower Forest Summer Wetland Floristic Survey*, unpublished report prepared for North Central Catchment Management Authority, Bendigo.

Australian Ecosystems. 2009b, *Gunbower Forest Spring Flora Survey*, unpublished report for the North Central Catchment Management Authority, Bendigo.

Australian Ecosystems. 2008a, *A Monitoring System for the Gunbower Forest*, revision of Francis Crome report (December 2004) for North Central Catchment Management Authority, Bendigo.

Australian Ecosystems. 2008b, *A Manual of Field Procedures for Monitoring in Gunbower Forest*, revision of Francis Crome report (December 2004) for North Central Catchment Management Authority, Bendigo.

Baker, M. C. 1979, 'Morphological correlates of habitat selection in a community of shorebirds', *Oikos*, vol. 33, pp. 121-126.


Balcombe, S., Arthington, A., Foster, N., Thoms, M., Wilson, G. and Bunn, S. 2006, 'Fish assemblages of an Australian dryland river: abundance, assemblage structure and recruitment patterns in the Warrego River, Murray-Darling Basin', *Marine and Freshwater Research*, vol. 57, pp. 619-633.

Baldwin, D. S. 1999, 'DOM and Dissolved P Leached from Fresh and 'Terrestrially'-Aged River Red Gum Leaves - Implications for Assessing River-Floodplain Interactions', *Freshwater Biology*, vol. 41, pp. 675-685.

Baldwin, D. S. and Mitchell, A. M. 2000, 'The effects of drying and re-flooding on the sediment and soil nutrient dynamics of lowland river-floodplain systems: a synthesis', *Regulated Rivers: Research and Management*, vol. 16, pp. 457-467.

Baumgartner, L. J., Boys, C. A., Stuart, I. G. and Zampatti, B. P. 2010, 'Evaluating migratory fish behaviour and fishway performance: testing a combined assessment methodology', *Australian Journal of Zoology*, vol. 58, pp. 154-164.

Baumgartner, L. J., Conallin, J., Wooden, I., Campbell, B., Gee, R., Robinson, W. A. and Mallen-Cooper, M. 2013, 'Using flow guilds of freshwater fish in an adaptive management framework to simplify environmental flow delivery for semi-arid riverine systems', *Fish and Fisheries*, Blackwell Publishing Ltd.



Baxter, C. V., Fausch, K. D. and Saunders, W. C. 2005, 'Tangled webs: reciprocal flows of invertebrate prey link streams and riparian zones', *Freshwater Biology*, vol. 50, pp. 201-220.

Bennetts, K. 2014, *Preliminary Vegetation Assessment of Benwell and Guttrum Forests*, report prepared for the North Central Catchment Management Authority, Fire, Flood and Flora, Victoria.

Bennetts, K. & Jolly, K. 2013, *Gunbower Forest Sentinel Wetland and Understorey Survey autumn 2013*, unpublished Technical Report for the North Central Catchment Management Authority, Fire, Flood and Flora, Victoria.

Biosis. 2014, *Flora and Fauna Assessment of the Gunbower National Park and Guttrum and Benwell State Forests*, Report for North Central Catchment Management Authority.

Blanch, S.J. and Brock, M.A. 1994, 'Effects of grazing and depth on two wetland plant species', *Australian Journal of Marine and Freshwater Research*, vol. 45, pp. 1387-94.

Blanch, S.J., Ganf, G.G. and Walker, K.F. 1998, 'Growth and recruitment in *Vallisneria americana* as related to average irradiance in the water column', *Aquatic Botany*, vol. 61, pp. 1810-205.

Bormans, M., Maier, H., Burch, M. and Baker, P. 1997, 'Temperature stratification in the lower River Murray, Australia: implication for cyanobacterial bloom development', *Marine and Freshwater Research*, vol. 48, pp. 647-654

Boulton, A. J. and Brock, M.A. 1991, 'Macroinvertebrate assemblages in floodplain habitats of the lower River Murray, South Australia', *Regulated Rivers: Research and Management*, vol. 6, pp. 183-201.

Brandis, K., Nairn, L., Porter, J. and Kingsford, R. T. 2009, *Preliminary assessment for the environmental water requirements of waterbird species in the Murray Darling*

Basin, June 2009, University of New South Wales, Sydney.

Bren, L. J. 1987, 'The duration of inundation in a flooding river red gum forest', *Australian Forest Research*, vol. 17, pp. 191-202.

Bren, L. J. 1988, 'Flooding characteristics of a riparian red gum forest', *Australian Forest Research*, vol. 51, pp. 57-62.

Bren, L. J. and Gibbs, N. L. 1986, 'Relationships between flood frequency, vegetation and topography in a River Red Gum forest', *Australian Forest Research*, vol. 16, pp. 357-370.

Briggs, S. V. 1990, Waterbirds, pages 337-344 in N. Mackay and D. Eastburn, editors, *The Murray*, Murray-Darling Basin Commission, Canberra.


Briggs, S.V. and Maher, M.T. 1985, 'Limnological studies of waterfowl habitat in south- western New South Wales. II. Aquatic macrophyte productivity', *Australian Journal of Marine and Freshwater Research*, vol. 36, pp. 707-15.

Briggs, S. V. and Maher, M. T. 1983, 'Litter fall and leaf decomposition in a River Red Gum (*Eucalyptus camaldulensis*) swamp', *Australian Journal of Botany*, vol. 31, pp. 307-316.

Briggs, S. A., Thornton, S. A. and Lawler, W. G. 1997, 'Relationship between hydrological control of river red gum wetlands and waterbird breeding', *Emu*, vol. 97, pp. 31-42.

Brookes, J., Burch, M., Wallace, T. and Baldwin, D. 2007, *Risk Assessment of Cyanobacteria and Blackwater Events in Chowilla Floodplain*, Report to the South Australian Murray-Darling Natural Resources Management Board.

Broome, L. S. and Jarman, P. J. 1983, 'Waterbirds on natural and artificial waterbodies in the Namoi Valley, New South Wales', *Emu*, vol. 83, pp. 99-104.



Butler, R. W. 1994, 'Population regulation of wading Ciconiiform birds', *Colonial Waterbirds*, vol. 17, pp. 189-199.

Cadwallader, P.L. 1977, *J. O. Langtry's 1949-50 Murray River Investigations*, Fisheries and Wildlife Paper, Victoria.

Chesterfield, E. A. 1986, 'Changes in the vegetation of the river red gum forest at Barmah, Victoria', *Australian Forestry*, vol. 49, pp. 4-15.

Colwell, M. A. and Taft, O. W. 2000, 'Waterbird communities in managed wetlands of varying water depth', *Waterbirds*, vol. 23, pp. 45-55.

Crome, F. 2004, *A Manual of Field Procedures for Monitoring in Gunbower Forest*, revised by Australian Ecosystems in 2008.

Crome, F. H. J. 1988, 'To drain or not to drain? - Intermittent swamp drainage and waterbird breeding', *Emu*, vol. 88, pp. 243-248.

Crook, D. 2004, 'Is the home range concept compatible with the movements of two species of lowland river fish?', *Journal of Animal Ecology*, vol. 73, pp. 353-366.

Davison, W. 1993, 'Iron and Manganese in lakes', *Earth-Science Reviews*, vol. 34, pp. 119-163.

Dexter, B. D. 1978, 'Silviculture of the River Red Gum forests of the central Murray floodplain', *Proceedings of the Royal Society of Victoria*, vol. 90, pp. 175-194.

DSE. 2003, *Gunbower Forest Ramsar Site Strategic Management Plan*, Department of Sustainability and Environment, Victoria.

Eaton, A. D., Clesceri, L.S. and Greenberg, A. E. 1995, *Standard Methods for the Examination of Water and Wastewater*, American Public Health Association: Washington, DC.

Ebner, B., Scholz, O. and Gawne, B. 2009, 'Golden perch *Macquaria ambigua* are flexible spawners in the Darling River, Australia', *New Zealand Journal of Marine and Freshwater Research*, vol. 43, issue 2, pp. 571-578.

Ecological Associates. 2014, *Preliminary Modelling Advice – draft report*, prepared for North Central Catchment Management Authority, Huntly.

Ecological Associates. 2013, *The Ecological Justification for Works and Measures for the Guttrum and Benwell State Forests*, Ecological Associates report AA019-1-A prepared for North Central Catchment Management Authority, Huntly.

Ecological Associates. 2010, *Gunbower Forest Hipwell Road Channel Ecological Benefit and Risk Analysis – Final Report*, Ecological Associates report AA-015-2-B prepared for North Central Catchment Management Authority, Huntly.


Ecological Associates. 2009, *Upper Forest Channel Risk Assessment*, Ecological Associates report AA014-1-B, unpublished report prepared for North Central Catchment Management Authority, Bendigo.

Ecological Associates. 2003, *Flooding Enhancement of Gunbower Forest – Investigation of Priority Options – Part A*, unpublished report prepared for North Central Catchment Management Authority, Bendigo.

Erskine, W.D., Saynor, M. J., Erskine, L., Evans, K. G., Moliere, D. R. 2005, 'A preliminary typology of Australian tropical rivers and implications for fish community ecology', *Marine and Freshwater Research*, vol. 56, pp. 253-267.

Fairbairn, S. E. and Dinsmore, J. J. 2001, 'Local and landscape-level influences on wetland bird communities of the prairie pothole region of Iowa, USA', *Wetlands*, vol. 21, pp. 41-47.

Feminella, J. W., Lockaby, B. G. and Schoonover, J. E. 2003, 'Land use change and stream signatures: effects



of urbanisation on stream biogeochemistry and biodiversity in catchments of western Georgia, USA', in *Proceedings of Symposium on Urbanisation and Stream Ecology*.

Fitzsimons, J. A., Peake, P., Frood, D., Mitchell, M., Withers, N., White, M. and Webster, R. 2011, 'Flooding requirements for biodiversity values along the Victorian floodplain of the Murray Valley', *The Victorian Field Naturalist*, vol. 28, issue 2, pp. 48-85.

Frood, D. 2012, *Water and Salinity Regime and Depth Preferences for Victorian Wetland Ecological Vegetation Classes*, Unpublished report prepared for the Department of Sustainability and Environment, Melbourne.

Furphy, G. 2006, *Threatened Grassy Vegetation Communities of the Goulburn Broken Catchment - Literature Review*, Department of Sustainability and Environment, Victoria.

Gawlik, D. E. 2002, 'The effects of prey availability on the numerical response of wading birds', *Ecological Monographs*, vol. 72, pp. 329-346.

Gawne, B. and Scholz, O. 2006, 'Synthesis of a new conceptual model to facilitate management of ephemeral deflation basin lakes', *Lakes and Reservoirs: Research and Management*, vol. 11, pp. 177-188.

Gehrke, P.C. 1997, 'Differences in composition and structure of fish communities associated with flow regulation in New South Wales Rivers', in *Fish and Rivers in Stress: The New South Wales Rivers Survey* (eds J. H. Harris and P. C. Gehrke), NSW Fisheries Office of Conservation and the Cooperative Research Centre for Freshwater Ecology, Sydney.

Gehrke, P. C. 1988, 'Response-surface analysis of teleost cardiorespiratory responses to temperature and dissolved oxygen', *Comparative Biochemistry and Physiology*, vol. 89, pp. 587-592.

Gehrke, P. C., Revell, M. B. and Philbey, A. W. 1993, 'Effects of river red gum, *Eucalyptus camaldulensis*, litter on golden perch, *Macquaria ambigua*', *Journal of Fish Biology*, vol. 43, pp. 265-279.

George, A. K. 2004, *Eucalypt Regeneration on the Lower Murray Floodplain, South Australia*, Thesis submitted in fulfilment of the requirements for the degree of Doctor of Philosophy, Adelaide: School of Earth and Environmental Sciences, The University of Adelaide.

George, A. K., Walker, K. F. & Lewis, M. M. 2005, 'Population status of eucalypt trees on the River Murray floodplain', *River Research and Applications*, vol. 21, pp. 271-282.


Gilligan, D. and Schiller, C. B. 2003, *Downstream Transport of Larval and Juvenile Fish in the River Murray*, NSW Fisheries Report Series No. 50, NSW Fisheries, Sydney, Australia.

Gilligan, D.M., Vey, A. and Asmus, M. 2009, *Identifying Drought Refuges in the Wakool System and Assessing Status of Fish Populations and Water Quality Before, During and After the Provision of Environmental, Stock and Domestic Flows*.

Hargeby, A., Andersson, G., Blindow, I. and Johansson, S. 1994, 'Trophic web structure in a shallow eutrophic lake during a dominance shift from phytoplankton to submerged macrophytes', *Hydrobiologia*, vol. 279/280, pp. 83-90.

Harris, J. H. and Gehrke, P. C. (eds). 1997, *Fish and Rivers in Stress: the NSW Rivers Survey*, NSW Fisheries Office of Conservation and the Cooperative Research Centre for Freshwater Ecology, Cronulla.

Harris, J. H., Thorncraft, G. A. and Wem, P. 1998, 'Evaluation of rock-ramp fishways in Australia', in M. Jungwirth, S. Schmutz and S. Weiss (eds) *Fish Migration and Fish Bypasses*, Oxford: Fishing News Books.



Heinrich, P. A. 1990, *The Ecophysiology of Riparian River Red Gum (Eucalyptus camaldulensis)*, Final Report AWRAC W86/13, Forestry Section, University of Melbourne, Melbourne.

Horrocks, G. F. B., Brown, G. W., Earl, G. E., Griffiths, R. C. and Williams, L. M. (unpublished), *Flora and Fauna of Gunbower Island, Northern Victoria: Ecological Survey*, Report No. 28, Department of Conservation, Forests and Lands, Melbourne.

Howitt, J., Baldwin, D. S., Rees, G. N. and Williams, J. 2007, 'Modelling blackwater: predicting water quality during flooding of lowland river forests', *Ecological Modelling*, vol. 203, pp. 229-242.

Humphries, P. 2005, 'Spawning time and early life history of Murray cod, *Maccullochella peelii peelii* (Mitchell) in an Australian River', *Environmental Biology of Fishes*, vol. 72, pp. 393-407.

Humphries, P., King, A. and Koehn, J. 1999, 'Fish, flows and flood plains: links between freshwater fishes and their environment in the Murray-Darling River system, Australia', *Environmental Biology of Fishes*, vol. 56, issue 1, pp. 129-151.

Humphries, P. and Potter, I. C. 1993, 'Relationship between the habitat and diet of three species of atherinids and three species of gobies in a temperate Australian estuary', *Marine Biology*, vol. 116, pp. 193-204.

Hutchison, M. et al. 2008, *Mesoscale Movements of Small and Medium-sized Fish in the Murray-Darling Basin*, Murray-Darling Basin Commission, Canberra.

Isola, C. R., Colwell, M. A., Taft, O. E. and Safran, R. J. 2000, 'Interspecific differences in habitat use of shorebirds and waterfowl foraging in managed wetlands of California's San Joaquin Valley', *Waterbirds*, vol. 23, pp. 196-203.

Jensen, A. E., Walker, K. F. and Paton, D. C. 2008, 'The role of seedbanks in restoration of floodplain

woodlands', *River Research and Applications*, vol. 24, pp. 632-649.

Johns, C. et al. 2009, *Literature Review and Experimental Design to Address Retaining Floodwater on Floodplains and Flow Enhancement Hypotheses Relevant to Native Tree Species*, Report prepared for the Murray-Darling Basin Authority by The Murray-Darling Freshwater Research Centre, Wodonga.

Jones, M. J. 2006, *Effects of Environmental Flow Allocations on the Lateral Movements of Native Fish in the Barmah-Millewa Forest*, Final report to the Murray-Darling Basin Authority, Arthur Rylah Institute for Environmental Research.

Jones, M. J. and Stuart, I.G. 2008, 'Regulated floodplains – a trap for unwary fish', *Fisheries Management and Ecology*, vol. 15, pp. 71-79.


Jones, M. and Stuart, I. 2008b, 'Lateral movement of common carp (*Cyprinus carpio*) in a large lowland river and floodplain', *Ecology of Freshwater Fish*, vol. 18, pp. 72-82.

Jones, M. and Stuart, I. G. 2004, 'Impacts of flow regulation structures on fish in the Barmah- Millewa Forest', *Freshwater Ecology*, Arthur Rylah Institute for Environmental Research.

Junk, W. J., Bayley, P. B. and Sparks, R. E. 1989, 'The flood pulse concept in river-floodplain systems', *Canadian Special Publication of Fisheries and Aquatic Sciences*, vol. 106, pp. 110-127.

Kerezszy, A., Balcombe, S. R., Arthington, A. H. and Bunn, S. E. 2011, 'Continuous recruitment underpins fish persistence in the arid rivers of far-western Queensland, Australia', *Marine and Freshwater Research*, vol. 62, issue 10, pp. 1178-1190.

King, A. J., Beesley, L., Petrie, R., Nielsen, D., Mahoney, J. and Tonkin, Z. 2011, *Monitoring Fish Recruitment, water Quality and Secondary Production in Barmah-*



Millewa Forest, 2010/11, Arthur Rylah Institute for Environmental Research, unpublished client report, Department of Sustainability and Environment, Heidelberg, Victoria.

King, A., Tonkin, Z. and Mahoney, J., 2009, 'Environmental flow enhances native fish spawning and recruitment in the Murray River, Australia', *River Research and Applications*, pp. 1205-1218.

King, A. J., Tonkin, Z. and Mahoney, J. 2007, *Assessing the Effectiveness of Environmental Flows on Fish Recruitment in Barmah-Millewa Forest*, Report to Murray-Darling Basin Commission, Freshwater Ecology, Arthur Rylah Institute for Environmental Research, Department of Sustainability and Environment.

King, A. J., Humphries, P. and Lake, P. S. 2003, 'Fish recruitment on floodplains: the roles of patterns of flooding and life history characteristics', *Canadian Journal of Fisheries and Aquatic Sciences*, vol. 60, pp. 773-786.

Kingsford, R. T. 1998, Management of wetlands for waterbirds in W. D. Williams, editor, *Wetlands in a Dry Land: Understanding for Management*, Environment Australia, Canberra.

Kingsford, R. T. and Auld, K. M. 2005, 'Waterbird breeding and environmental flow management in the Macquarie Marshes, arid Australia', *River Research and Applications*, vol. 21, pp. 187-200.

Kingsford, R. T. and Norman, F. I. 2002, Australian waterbirds - products of the continent's ecology', *Emu*, vol. 102, pp. 47-69.

Kingsford, R. T. and Porter, J. L. 1994, 'Waterbirds on an adjacent freshwater lake and salt lake in arid Australia', *Biological Conservation*, vol. 65, pp. 141-151.

Kingston, R. T., Jenkins, K. M. and Porter, J. L. 2004, 'Imposed hydrological stability on lakes in arid Australia

and effects on waterbirds', *Ecology*, vol. 85, pp. 2478-2492.

Koehn, J. 2009, 'Multi-scale habitat selection by Murray cod *Maccullochella peelii peelii* in two lowland rivers', *Journal of Fish Biology*, vol. 75, issue 1, pp. 113-129.

Koehn, J. D. 2006, *The Ecology and Conservation of Murray Cod (Maccullochella peelii peelii)*, PhD thesis, University of Melbourne.

Koehn, J., McKenzie, J., O'Mahony, D., Nicol, S., O'Connor, J. and O'Connor, W. 2009, 'Movements of Murray cod (*Maccullochella peelii peelii*) in a large Australian lowland river', *Ecology of Freshwater Fish*, vol. 18, issue 4, pp. 594-602.


Koehn, J. D., Brumley, A. and Gehrke, P. C. 2000, *Managing the Impacts of Carp*, Bureau of Rural Sciences, Department of Agriculture, Fisheries and Forestry, Canberra, Australia.

Koehn, J., Nicol, S., McKenzie, J., Lieschke, J., Lyon, J. and Pomorin, K. 2008, 'Spatial ecology of an endangered native Australian Percichthyid fish, the trout cod *Maccullochella macquariensis*', *Endangered Species Research*, vol. 4, issues 1-2, pp. 219-225.

Koehn, J. and Nicol, S. 1998, 'Habitat and movement requirements of fish,' in R. J. Banens and R. Lehane (eds). Proceedings of the 1996 Riverine Environment Research Forum, Murray Darling Basin Commission, Canberra.

Koehn, J. D. and O'connor, W. G. 1990, *Biological Information for Management of Native Freshwater Fish in Victoria*, Department of Conservation and Environment Freshwater Fish Management Branch Arthur Rylah Institute for Environmental Research, Melbourne.

Lake, J. S. 1967, 'Rearing experiments with five species of Australian freshwater fishes. I. Inducement to



spawning', *Australian Journal of Marine and Freshwater Research*, vol. 18, pp. 155–173.

Lawrence, I. and Breen, P. 1998, *Design Guidelines: Stormwater Pollution Control Ponds and Wetlands*, The Cooperative Research Centre for Freshwater Ecology, Canberra.

Laws, E. A. 1993, *Aquatic Pollution: An Introductory Text*, Wiley and Sons Ltd, New York.

Leck, M. A. & Brock, M. A. 2000, 'Ecological and evolutionary trends in wetlands: Evidence from seeds and seed banks in New South Wales, Australia and New Jersey, USA', *Plant Species Biology*. vol. 15, pp. 97–112.

Leigh, S. J. and Zampatti, B. 2011, *Movement and Spawning of Murray Cod (*Maccullochella peelii*) and Golden Perch (*Macquaria ambigua*) in Response to a Small-scale Flow Manipulation in the Chowilla Anabranch System*, SARDI publication No. F2010/000646-1.

Leitch, C. 1989, *Towards a Strategy for Managing the Flooding of Barmah Forest*, Department of Conservation Forests and Lands, Benalla, Victoria.

Leslie, D. J. 2001, 'Effect of river management on colonially-nesting waterbirds in the Barmah-Millewa forest, south-eastern Australia', *Regulated Rivers: Research and Management*, vol. 17, pp. 21-36.

Lintermans, M. 2009, *Fishes of the Murray-Darling Basin: An introductory Guide*, Murray-Darling Basin Authority, Canberra.

Lintermans, M. 2007, *Fishes of the Murray-Darling Basin: An introductory Guide*, Murray-Darling Basin Commission, Canberra.

Lovvorn, J. R. 1994, 'Biomechanics and foraging profitability - an approach to assessing trophic needs and impacts in foraging ducks' *Hydrobiologia*, vol. 280, pp. 223-233.

Loyn, R. H., Lumsden, L. F. & Ward, K. A. 2002, 'Vertebrate fauna of Barmah Forest, a large forest of River Red Gum *Eucalyptus camaldulensis* on the floodplain of the Murray River', *The Victorian Naturalist*, vol. 119, version 3, pp. 114-123.

Lyon, J., Stuart, I., Ramsey, D. and O'Mahony, J. 2010, 'The effect of water level on lateral movements of fish between river and off-channel habitats and implications for management', *Marine and Freshwater Research*, vol. 61, pp. 271-278.

Mallen-Cooper, M. 1999, 'Developing fishways for non-salmonid fishes; a case study from the River Murray in Australia', in *Innovations in Fish Passage Technology*, Odeh, M. (ed.). American Fisheries Society: Bethesda.


Mallen-Cooper, M., Koehn, J., King, A., Stuart, I. and Zampatti, B. 2007, *Risk Assessment of the Proposed Chowilla Regulator and Managed Floodplain Inundations on Fish*, Report to DLWBC.

Mallen-Cooper, M. and Stuart, I. G. 2003, 'Age, growth, and non-flood recruitment of two potamodromous fishes in a large semi-arid/temperate river system', *River Research and Applications*, vol. 19, pp. 697–719.

Mallen-Cooper, M., Stuart, I. G. & Sharpe, C. 2014, *The Gunbower – Lower Loddon Native Fish Recovery Plan*, report produced for North Central Catchment Management Authority, Bendigo.

Marcar, N. E. 1993, 'Waterlogging modifies growth water use and ion concentration in seedlings of salt-treated *Eucalyptus camaldulensis*, *E. tereticornis*, *E. robusta* and *E. Globulus*', *Australian Journal of Plant Physiology*, vol. 20 pp. 1-13.

Martinova, M. V. 1993, 'Nitrogen and phosphorus compounds in bottom sediments: mechanisms of accumulation, transformation and release', *Hydrobiologia*, vol. 252, pp. 1-22.



McDougall, A. and Timms, B. 2001, 'The influence of turbid waters on waterbird numbers and diversity: a comparison of Lakes Yumberarra and Karatta, Currawinya National Park, south-west Queensland', *Corella*, vol. 25, pp. 25-31.

McDowall, R. M. (ed.) 1996, *Freshwater Fishes of South-Eastern Australia*, Reed Books, Sydney.

McEvoy, P. 1992. *Ecophysiological Comparisons Between Eucalyptus camaldulensis Dehnh., E. largiflorens F. Muell. and E. microcarpa (Maiden) Maiden on the River Murray Floodplain*, Unpublished M. Forest Science, University Melbourne, Melbourne.

McMaster, D. and Bond, N. 2008, 'A field and experimental study on the tolerances of fish to *Eucalyptus camaldulensis* leachate and low dissolved oxygen concentrations', *Marine and Freshwater Research*, vol. 59, pp. 177-185.

McNeil, D., Wilson, P., Hartwell, D. and Pellizzari, M. 2008, *Olive Perchlet (Ambassis agassizii) in the Lachlan River: Population Status and Sustainability in the Lake Brewster Region*, a report submitted to the Lachlan Catchment Management Authority, SARDI Aquatic Sciences Publication No F2008/000846-1, SARDI Research Report Series No 309.

Meredith, S. 2006, 'Flows, fish and wetlands', in Ward, T. M., Geddes, M. C. and Mayfield, S. *National Symposium on Ecosystem Research and Management of Fish and Fisheries*, Australian Society for Fish Biology Symposium Proceedings, Adelaide, South Australia, September 2004.

Morin, J. and Morse, J. W. 1999, 'Ammonium release from resuspended sediments in the Laguna Madre estuary', *Marine Chemistry*, vol. 65, pp. 97-110.

Mortimer, C. H. 1941, 'The exchange of dissolved substances between mud and water in lakes (parts I and II)', *Journal of Ecology*, vol. 29, pp. 280-329.

Murray Darling Basin Authority. 2012, *Gunbower Forest Environmental Water Management Plan*, MDBA, Canberra.

Murray Darling Basin Authority. 2010, *Additional Hydraulic Modelling of TLM Interventions in Gunbower Forest Victoria*, unpublished report, Murray Darling Basin Authority, Canberra.

North Central CMA. 2014a, *Flooding Enhancement of Gunbower Forest Ecological Watering Guide*, Report prepared as a requirement of The Living Murray program.

North Central CMA. 2014b, *Seasonal Watering Proposal for Gunbower Forest and Gunbower Creek 2014-15*, draft, North Central CMA, Huntly.

North Central CMA. 2013, *Flooding Enhancement of Gunbower Forest Interim Operating Plan*, draft prepared as a requirement of The Living Murray Program.


North Central CMA. 2012, *Pig Swamp Environmental Watering Plan*, Report prepared for the Goulburn Murray Water Connections Project, North Central CMA, Huntly.

North Central CMA. 2010, *Flooding Enhancement of Gunbower Forest Project Investment Proposal – Gunbower-Koondrook-Perricoota Forests*, proposal to the Murray Darling Basin Authority.

North Central CMA. 2009, *Gunbower Waterbird Breeding Requirements Workshop – Summary Paper*, draft, unpublished report prepared by Chatfield, A. & Tranter, M., Bendigo.

O'Connell, M., Baldwin, D. S., Rees, G. and Robertson, A. 2000, 'Release and bioavailability of dissolved organic matter from floodplain litter: Influence of origin and oxygen levels', *Freshwater Biology*, vol. 45, pp. 333-342.

O'Connor, J., King, A., Tonkin, Z., Morrongiello, J. and Todd, C. 2008, *Fish in the Murray Valley and Torrumbarry Irrigation Areas*, Arthur Rylah Institute for



Environmental Research Technical Report Series No. 176, Department of Sustainability and Environment, Heidelberg, Victoria.

O'Connor, J. P., O'Mahony, D. J. and O'Mahony, J. M. 2005, 'Movements of *Macquaria ambigua*, in the River Murray, south-eastern Australia', *Journal of Fish Biology*, vol. 66, pp. 392–403.

O'Connor, P. G. & Ward, K. A. 2003, *Waterbird Monitoring in Barmah Forest 2002-2003*, Department of Sustainability and Environment, Tatura, Victoria.

Parsons, W. T. and Cuthbertson, E. G. 1992, *Noxious Weeds of Australia*, Inkata Press, Melbourne.

PIRVic. 2007, *Report of a Fish Survey of the Gunbower Creek*, Progress report to NCCMA, Huntly.

Pusey, B., Kennard, M. and Arthington, A. 2004, *Freshwater Fishes of North-Eastern Australia*, CSIRO Publishing, Collingwood, Australia.

Rehwinkel, R. and Sharpe, C. 2009, *Gunbower Forest Fish Monitoring Surveys*, MDFRC Report to NCCMA.

Reynolds, L. F. 1983, 'Migration patterns of five fish species in the Murray-Darling River system', *Australian Journal of Marine and Freshwater Research*, vol. 34, pp. 857–871.

Richardson, A., Meredith, S., Conallin, A. and Sharpe, C. 2005, *Status of the Gunbower Island Fish Community, June 2005: including Recommendations for Future Monitoring*, Report prepared for North Central Catchment Management Authority, November.

Roberts, J. and Marston, F. 2000, *Water Regime of Wetland and Floodplain Plants in the Murray-Darling Basin: a Source Book of Ecological Knowledge*, CSIRO Technical Report 30/00, CSIRO Land and Water, Canberra.

Roberts, J. & Marston, F. 2011, *Water regime for wetland and floodplain plants: a source book for the*

Murray–Darling Basin, National Water Commission, Canberra.

Robertson, A. I., Bacon, P. & Heagney, G. 2001, 'The responses of floodplain primary production to flood frequency and timing', *Journal of Applied Ecology*, vol. 38, pp. 126-136.

Rogers, K. (in prep.), *Water Regime Requirements for Selected Floodplain Wetland Plants at the Macquarie Marshes*, Department of Environment and Climate Change, Rivers and Wetlands Unit, Canberra.

Rogers, K. & Ralph, T. J. 2011, *Floodplain Wetland Biota in the Murray-Darling Basin Water and Habitat Requirements*, CSIRO Publishing, Collingwood.

Rolls, R. and Wilson, G. 2010, 'Spatial and temporal patterns in fish assemblages following an artificially extended floodplain inundation event, northern Murray-Darling Basin, Australia', *Environmental Management*, vol. 45, issue 4, pp. 822-833.


Rowland, S. J. 1983, 'Spawning of the Australian freshwater fish Murray cod, *Maccullochella peelii* (Mitchell), in earthen ponds', *Journal of Fish Biology*, vol. 23, pp. 525–534.

Saddler, S. and O'Mahony, J. 2007, *Protection and Enhancement of Murray Cod Populations*, Arthur Rylah Institute for Environmental Research, Report prepared for the Living Murray Program, Murray-Darling Basin Commission.

Safran, R. J., Colwell, M. A. Isola, C. R. and Taft, O. E. 2000, 'Foraging site selection by nonbreeding white-faced ibis', *The Condor*, vol. 102, pp. 211-215.

Scott, A. 1997, *Relationship between waterbird ecology and river flows in The Murray-Darling Basin*, CSIRO Technical report 5/97, Canberra.

Serena and Williams. 2013, *Water-related Requirements for Platypus Management in the Gunbower Lagoons*, A



Report to the North Central CMA and Goulburn-Murray Water.

Shang, E. H. H. and Wu, R. S. S. 2004, 'Aquatic hypoxia is a teratogen and affects fish embryonic development', *Environmental Science & Technology*, vol. 38, pp. 4763 - 4767.

Sharpe, C.P. 2011, *Spawning and Recruitment Ecology of Golden Perch (Macquaria ambigua Richardson 1845) in the Murray and Darling Rivers*, PhD Thesis, Griffith School of Environment, Griffith University, Brisbane.

Sharpe, C. and Stuart, I. 2013, *Large Bodied Fish Hydrograph*, In presentation to the GEWAG 29th January 2013, Unpublished, Adapted from Baumgartner L., Conallin, J., Wooden, I., Campbell, B., Gee, R., Robinson, W. and Mallen-Cooper, M. 2013, 'Using flow guilds of freshwater fish in an adaptive management framework to simplify environmental flow delivery for semi-arid riverine systems', in *Fish and Fisheries*, Blackwell Publishing.

Sharpe, C. and Rehwinkel, R. 2009, *Status of Freshwater Catfish Populations in Gunbower Creek*, Preliminary progress report to NCCMA, MDFRC publication.

Sharpe, C., Campbell-Brown, S., Vilizzi, L. and Stanislawski, K. 2013, *Gunbower Island Annual Fish Surveys: 2013*, Report for the North Central Catchment Management Authority by CPS Environmental.

Sharpe, C., Campbell-Brown, S. and Vilizzi, L. 2012, *Gunbower Island Annual Fish Surveys: Autumn 2012*, Report for the North Central Catchment Management Authority, CPS Environmental Research, November 2012.

Simpson, K., Smith, J. N. M. and Kelsall, J. P. 1987, 'Correlates and consequences of coloniality in great blue herons', *Canadian Journal of Zoology*, vol. 65, pp. 572-577.

SKM. 2009, *Gunbower Forest Upper Forest Channel - Ecological Constraints Analysis*, Sinclair Knight Merz, Malvern, Victoria.

SKM. 2007, *Guttrum and Benwell State Forests Water Management Investigation – Water Management Options*, final report prepared for North Central CMA.

Slavich, P. G., Walker, G. R., Jolly, I. D., Hatton, T. J. and Dawes, W. R. 1999, 'Dynamics of *Eucalyptus largiflorens* growth and water use in response to modified watertable and flooding regimes on a saline floodplain', *Agricultural Water Management*, vol. 39, pp. 245-264.


Smith, B., Thwaites, L., Hillyard, K., Conallin, A. and Higham, J. 2008, 'MDBC Project MD746 Evaluation of Carp Exclusion Screens at Wetland Inlets: Existing Designs, Consideration of Ecological Effects & Conceptual Development of 'Optimised' Screens', in *Proceedings of the Murray-Darling Basin Commission Native Fish Strategy Forum 2007* (Ed. M. Lintermans).

Stuart, I. G. and Jones, M. 2006, 'Large, regulated forest floodplain is an ideal recruitment zone for non-native common carp (*Cyprinus carpio* L.)', *Marine and Freshwater Research*, vol. 57, pp. 333-347.

Stuart, I.G. and Mallen-Cooper, M. 1999, 'An assessment of the effectiveness of a vertical-slot fishway for non-salmonid fish at a tidal barrier on a large tropical/subtropical river', *Regulated Rivers: Research & Management*, vol. 15, pp. 575-590.

Stuart, I. G., Baumgartner, L. J. and Zampatti, B. P. 2008, 'Lock gates improve passage of small-bodied fish and crustaceans in a low gradient vertical-slot fishway', *Fisheries Management and Ecology*, vol. 15, pp. 241-248.

Stuart, I.G., Zampatti, B.P. and Baumgartner, L. J. 2008, 'Can a low gradient vertical-slot fishway provide passage for a lowland river fish assemblage?' *Marine and Freshwater Research*, vol. 59, pp. 332-346.



Taft, O. E., Colwell, M. A., Isola, C. R. and Safran, R. J. 2002, 'Waterbird responses to experimental drawdown: implications for the multispecific management of wetland mosaics', *Journal of Applied Ecology*, vol. 39, pp. 987-1001.

Turner, L. and Erskine, W. D. 2005, 'Variability in the development, persistence and breakdown of thermal, oxygen and salt stratification on regulated rivers of southeastern Australia', *River Research and Applications*, vol. 21, pp. 151-168.

URS. 2001, *Flooding Enhancement of Gunbower Forest: Scoping Study*, unpublished report prepared for North Central Catchment Management Authority, Bendigo.

VanRees-Siewert, K. L. and Dinsmore, J. J. 1996, 'Influences of wetland age on bird use of restored wetlands in Iowa', *Wetlands*, vol. 16, pp. 577-582.

VEWH. 2014, *Guidelines for Seasonal Watering Proposals 2014-15*, updated 8 January, Victorian Environmental Water Holder.

Wallace, T. A. 2008, *Water Quality Within Two Contrasting Wetlands at Chowilla Floodplain following Pondered Flooding*, Report to the SA Murray-Darling Basin Natural Resources Management Board, Murray-Darling Freshwater Research Centre, Mildura.

Wallace, T. A., Sharpe, C., Fraser, P., Rehwinkel, R. and Vilizzi, L. 2008, *The Impact of Drought on Water Quality and Fish Communities within Refuge Pools on the Lower Darling River*, A technical report prepared for the Lower Murray Darling Catchment Management Authority, Murray-Darling Freshwater Research Centre, Mildura.

Walsh, C. J. 2002, 'voiding going down the drain', in *Watershed*, Cooperative Research Centre for Catchment Hydrology.

Walsh, C. J., Sharpe, A. K., Breen, P. F. and Sonneman, J. A. 2001, 'Effects of urbanisation on streams of the Melbourne region, Victoria, Australia. 1. Benthic

macroinvertebrate communities', *Freshwater Biology*, vol. 46, pp. 535-551.

Ward, K. 2005, 'Water management in the changing Barmah-Millewa wetlands', *Proceedings of the Royal Society of Victoria*, vol. 117, issue 1, pp. 77-84.

Ward, K. 2002, *Monitoring Frog Response to Flooding in Barmah Forest 2001-02*, Final report prepared for the Barmah-Millewa Forum.

Ward, K. 2001, *Monitoring Frog Response to Flooding in Barmah Forest 2000-01*, Final report prepared for the Barmah-Millewa Forum.

Ward, K. 1996, *Flood Requirements of Vegetation for the Rehabilitation of Lake Mokoan*, Department of Natural Resources and Environment, Unpublished draft report.

Ward, P. 2009, *Monitoring Frog Response to Flooding in Gunbower Forest*, June sentinel study site monitoring report, prepared for the North Central CMA, Huntly.


Webster, R. 2004, *The Effect of Flooding on Waterbirds within Barmah-Millewa Forest - 1999-2003*, Ecosurveys Pty Ltd, Deniliquin.

Wu, R. S. S., Zhou, S. B., Randall, D. J., Woo, N. Y. S. and Lam, P. K. S. 2003, 'Aquatic hypoxia is an endocrine disruptor and impairs fish reproduction', *Environmental Science & Technology*, vol. 37, pp. 1137-1141.

Yen, S. and Myerscough, P.J. 1989, 'Co-existence of three species of amphibious plants in relation to spatial and temporal variation: Investigation of plant responses', *Australian Journal of Ecology*, vol. 14, pp. 305-318.

Young, W., Scott, A., Cuddy, S. and Rennie, B. 2003, *Murray Flow Assessment Tool: a Technical Description*, CSIRO Land and Water, Canberra.

Zampatti, B. and Leigh, S. 2013, 'Within-channel flow promotes spawning and recruitment of golden perch,




Macquaria ambigua ambigua- implications for environmental flow management in the River Murray, Australia', *Marine and Freshwater Research*, vol. 64, pp. 618-630.

Zampatti, B., Leigh, S., Nicol, J. and Weedon, J. 2006b, *2006 Progress Report for the Chowilla Fish and Aquatic Macrophyte Project*, Report to the Department of Land Water and Biodiversity Conservation.

Zampatti, B., Nicol, J., Leigh, S. and Bice, C. 2006a. *2005 Progress Report for the Chowilla Fish and Aquatic Macrophyte Project*, Report to the Department of Land Water and Biodiversity Conservation.

Zeffer, A., Johansson, L. C. and Marmebro, A. 2003, 'Functional correlation between habitat use and leg morphology in birds (Aves)', *Biological Journal of the Linnean Society*, vol. 79, pp. 461-484.



Appendix 1: Literature review of hydrological requirements

Watering risks to be managed

Avoid delivery that supplies shallow water over summer months, which may result in displacement of wetland flora through **excessive River Red Gum recruitment** (Ecological Associates 2010). Alternatively, ensure water is available for follow-up flooding to drown red gum saplings.


Several **aquatic weeds** have the potential to expand their distributions into Benwell Forest through propagules being transported in environmental water and suitable conditions for growth being provided under the environmental water regimes (Ecological Associates 2010). These species need to be monitored and management actions adopted where appropriate. High threat weeds include those with the potential to invade from the broader region (Alligator Weed (*Alternanthera philoxeroides*), Senegal Tea Plant (*Gymnocoronis spilanthoides*) and Cabomba (*Cabomba caroliniana*)). Some terrestrial weeds may also benefit from increased soil moisture provided through environmental watering including Bridal Creeper (*Asparagus asparagoides*), Blackberry (*Rubus fruticosus* spp. agg.), Horehound (*Marrubium vulgare*), Patterson's Curse (*Echium plantagineum*), Prickly Pear (*Opuntia* spp.), African Box-thorn (*Lycium ferocissimum*) and thistles (namely *Sonchus* spp. and *Cirsium vulgare*) (Ecological Associates 2010).

Ensure **contingency environmental water is available to support bird breeding events** that may initiate as a result of environmental water delivery to wetland systems. Bird breeding may be a primary goal of an environmental watering event if catchment conditions are appropriate (e.g. ideal climatic cues) or it may be an unintended (but desirable) ecological outcome. Either way, if a breeding event occurs, careful monitoring of the area of inundation and water depth around breeding sites (beneath nests and in foraging areas) will need to occur. Supplementary environmental water may be required, particularly early in the breeding cycle (August to October), to maintain the depth and area of inundation in key areas so that waterbirds do not abandon their nests and chicks before they fledge (Ecological Associates 2010).

Carp spawning and recruitment is likely to increase from delivering environmental water to Benwell Forest. Common Carp are found in the River Murray and they will use inundated floodplains to spawn on newly inundated vegetated ground, under a wide range of temperatures and time periods within the breeding season. Where possible, limit summer flooding (inundation after December) as this is highly advantageous to carp which have a longer breeding season than native fish (Ecological Associates 2010).

Native fish (in particular larvae) in the irrigation channel water may enter Benwell Forest during water delivery events. To **avoid fish being stranded** in Benwell Forest, ensure a sharp drop in water level is provided in late spring or summer (to provide fish with a cue to leave the forest) followed by a gradual ongoing decline in water level (to provide fish with opportunities to leave the floodplain while the wetland and riverine habitats are still connected) (Ecological Associates 2010).

Blackwater events occur when organic matter on the floodplain is inundated and the process of decay uses dissolved oxygen in the water much faster than it can be produced (through the air-water interface or plant photosynthesis). Low dissolved oxygen levels (below 4 mgO₂/L) can stress native fish and other aquatic animals, while very low dissolved oxygen (e.g. 0.5mgO₂/L anoxic conditions) can kill aquatic organisms. While blackwater is a natural phenomenon, it can be reduced by regular inundation (to avoid high loads of organic matter build up), avoiding inundation in summer months (dissolved oxygen declines as temperature increases) and minimising the creation of still or deep water environments that are prone to stratification (Ecological Associates 2010). Avoid delivering environmental water when there is a combination of lower than normal



flow in the River Murray and higher than average temperatures. For environmental watering post long, dry periods in Benwell Forest, deliver the flow when there is moderate to high River Murray flows to allow a full assessment of water quality impacts to determine the minimal safe flow for operation. Ensure contingency water is available in upstream storages for dilution purposes if water quality deteriorates.

Semi-permanent wetlands

Flood frequency and duration

Ecological Associates (2013) noted that water would normally be present in the semi-permanent wetlands of the Guttrum and Benwell Forests, but they are shallow and would dry out from time to time.

The semi-permanent wetlands across the two systems share a similar hydrology. Inflows commence between 17,000 to 23,000 ML/d, and under natural conditions occur approximately 9 out of 10 years with events lasting 3 to 6 months (interquartile range at 21,000 ML/d). The wetlands retain water on the flood recession to a depth of 0.5 to 0.7 m and would usually remain flooded during summer. The wetlands would often dry out in autumn but in wet years may remain flooded until the following winter. Reed Bed Swamp retains water to a greater depth of 0.85 m and is more likely to remain flooded throughout the year (Ecological Associates 2013).

Similar to the semi-permanent wetlands within Gunbower Island, these semi-permanent wetlands require a drying phase at least every two years to maintain their biological diversity (Ecological Associates 2010).


Example aquatic herbs that may be found in these wetlands include Swamp Lily (*Ottelia ovalifolia*) and Clove strip (*Ludwigia peploides* subsp. *montevidensis*). Swamp Lily is found where flooding lasts 2-6 months and Clove strip occurs where floods last 8-10 months (Ward 1996).

SKM (2007) suggested the semi-permanent wetlands should remain full for at least the length of the wetland plant growing season (spring through to the end of summer for submerged and emergent plants), so they can complete their lifecycle by flowering and contributing to the seed bank. Preferably, semi-permanent wetlands would remain full until the following growing season.

Timing and depth of flooding

Aquatic herblands are characterised by a diversity of small to large emergent herbs (i.e. Swamp Lily, Milfoils (*Myriophyllum* spp.), Clove-strip, and Star Fruit (*Damasonium minus*)) that emerge in shallow (often less than 50 centimetre) waters of semi-permanent wetlands, and at the seasonally inundated edge of permanent wetlands. It is a combination of wetting and drying rather than season that has been found to drive germination in similar wetland vegetation in Australia (Leck and Brock 2000). Furthermore wetland monitoring in Gunbower Forest post environmental flows suggests species diversity in this vegetation type is reduced with increasing water depth (Australian Ecosystems 2009). Roberts & Marston (2000) reported similar sensitivities of the characteristic species and that their germination from seed and propagules is highest following autumn and spring flooding. The persistence of aquatic herbland is hence potentially threatened by inappropriately timed or deep flooding, and at the other end of the scale shallow flooding prior to summer that promotes River Red Gum colonisation. River Red Gums in this context have the potential to shade and out compete for resources (Ecological Associates 2010).

In northern Victoria, Swamp Lily is found in shallow waters to 50 cm deep, where flooding occurs from winter to summer (Ward 1996). Clove-strip is found where flooding occurs in winter-summer, to depth of one metre (Ward 1996). Seeds germinate under water and on wet soil but require light (Yen and Myerscough 1989).



Germination is temperature sensitive, with no germination at 10°C compared with an optimum at 30°C: at this temperature, germination starts within a day of flooding and is completed in less than 5 days, but at higher and at lower temperatures, germination is delayed, and at 40°C success rate is halved (Yen and Myerscough 1989) (Roberts & Marston 2000).

River Red Gum with flood dependent understorey

Note: The below predominantly outlines supporting evidence for the hydrological requirements of River Red Gums (*Eucalyptus camaldulensis*). However, when ‘designing’ a water regime for a River Red Gum forest **the requirements of other species within the understorey need to be considered**. Understorey species have seasonal responses, which can lead to changes in community composition and in forest structure (Robertson et al. 2000, Bren 1987).

Cooling (2003) indicated that the following understorey species may be helpful in characterising the water regime for River Red Gum with flood dependent understorey - *Triglochin* spp., *Eleocharis acuta*, *Paspalidium jubiflorum*, *Alternanthera denticulata*, *Cynodon dactylon* var *pulchellus*, *Juncus subsecundus* and *Poa labillardierei* (Crome 2004). Ecological Associates (2013) also found these species within the Guttrum and Benwell forest understorey – *Carex tereticaulis*, *Juncus usitatus* combined with *Phragmites australis*, *Typha domingensis* and *Juncus ingens* in local depressions and seasonal, submerged aquatic macrophytes including *Eleocharis acuta* and *Alternanthera denticulata*. EVCs applicable to this vegetation community include Grassy Riverine Forest (EVC 106), Riverine Swamp Forest (EVC 814) and Sedgy Riverine Forest (EVC 816) (K. Bennetts, pers. comm. August 2014).


Flood frequency and duration

Red gum with flood-dependent understorey occurs in Benwell Forest in areas that have a low flooding threshold but do not retain deep water when flood water recedes. At Benwell Forest flooding commences at flows exceeding 18,000 ML/d. Flooding of the Red Gum Forest with flood dependent understorey is largely complete at flows of 26,000 ML/d. Under natural conditions the forest would have been flooded in approximately 8 years out of 10 for 2 to 5 months (interquartile range at 23,000 ML/d) with flooding typically occurring between June and December.

The minimum duration of flooding for this vegetation type is 60 days according to Cunningham et al. (2009). This would include a flow peak and progressive drawdown.

Colonial waterbirds require suitable places to build their nests in order to breed successfully. For colonial stick-nesting waterbirds dense vegetation is essential for breeding (Kingsford and Norman 2002). Breeding in these species is dependent on areas of living River Red Gum that are flooded for at least four months (Briggs et al. 1997).

Current flood frequency for most of the River Red Gum forest at Barmah is about 6-8 years in every 10, on average (Bren and Gibbs 1986). Historical analyses of flood records shows that forest trees used to (prior to Hume Dam) experience inundation for 1-7 months and that this occurred primarily in winter- spring. There is increasing evidence that duration is as important as frequency, in terms of whole-forest growth responses. For trees that have been through a dry phase, frequent short floods and longer floods both reduce water stress and hence result in greater growth (Roberts & Marston 2000).



At Barmah, River Red Gums are known to have tolerated relatively long periods of continuous flooding, estimated as 24 months (Bren 1987). This has happened at least twice in very wet periods, once in the mid-1950s and once in the mid-1970s. This estimate of 24 months is consistent with several field observations of about 2-4 years of continuous flooding, before trees show signs of stress. However, this would apply to the wettest River Red Gum communities (D. Flood, pers. comm. August 2014). Trees behind Hay Weir apparently survived 3- 4 years continuous inundation (Bren 1987); River Red Gums at Murrumbidgee Swamp on the Lachlan River which were flooded continuously between 1974 and 1977 showed no signs of stress worth reporting (Briggs and Maher 1983) in that time; four wet years killed off some low-lying trees in Barmah Forest (Chesterfield 1986). Variations in these estimates and in field observations are due to differences and patchiness in soil properties, air spaces and in root respiratory demands (Roberts & Marston 2000).


River Red Gum seedlings are sensitive to prolonged inundation or high temperatures over summer and/or frost during winter (Ecological Associates 2010).

Regarding understorey species, the following inundation frequency and duration requirements have been documented:

- *Lobelia concolor* occurs in northern Victoria where flooding lasts 1-3 months (Ward 1996).
- *Triglochin procerum* has an optimum duration in northern Victoria of 6 months, but can tolerate 1-8 months (Roberts and Marston 2000).
- *Eleocharis acuta* in northern Victoria has an optimum flood duration of 8 months, but can tolerate 3-10 months of flooding (Ward 1996).
- *Paspalidium jubiflorum* (Warrego Summer Grass) is found where flooding occurs for 2-4 months (70-140 days) per year (not necessarily consecutive) (Roberts and Marston 2000).
- *Carex tereticaulis* tolerates flooding of 1-4 months in northern Victoria, but optimum duration is 2 months (Roberts and Marston 2000).
- *Phragmites australis* does not require flooding to survive, only adequate water. On the banks of the River Murray in South Australia, *Phragmites* occurs where flooding occurs for 80-225 days per year, not necessarily consecutive days (Blanch et al. 1999).
- *Typha domingensis* grows where the water regime ranges from permanently wet to seasonally or periodically dry. It can tolerate dry conditions for short periods (3-4 months in summer-autumn) once the growing season is over, without loss of vigour. The rhizome can survive dry conditions for even longer, possibly a few years, if protected from desiccation by being deep (0.5 m) within heavy clay. On the banks of the River Murray, South Australia, *Typha domingensis* is found in permanently flooded or moist sites where flooding occurs for 360-365 days per year (Roberts and Marston 2000).
- *Juncus ingens* occurs in northern Victoria, where flooding is in winter–spring and lasts for 6-11 months. However, the optimum duration is 9 months (Ward 1996).

Timing of flooding

Flooding of River Red Gum with flood dependent understorey occurs mainly in winter and spring. Flooding in spring provides a shallow, productive habitat for aquatic plants to develop and in which small fish reproduce, and breeding waterbirds and large fish find prey. The recession of water before summer provides germination



opportunities for a number of understorey species in the damp soil. A number of perennial species, such as Warrego Summer Grass and *Lobelia concolor*, grow on the forest floor over summer. Sustained flooding through summer prevents the establishment of these species, resulting in lower vegetation cover through autumn (Ecological Associates 2010).

Flood timing affects germination success for River Red Gum trees. For example, winter floods with winter recessions usually provide unfavourable water and air temperatures for seeds. Spring-summer floods followed by summer recession provide suitable germination conditions but subsequent heat and water stress can cause massive seedling mortality. Regeneration is optimised if flood recession is in spring-early summer, as this results in 'prolific' germination (Dexter 1978).

Production in River Red Gum trees was found to be higher where summer floods or spring and summer floods were received. Production was lower where only spring floods, or no floods occurred (Robertson et al. 2001).

Tree growth (i.e. wood production) in River Red Gums is greatest when flooded under warm conditions such as summer (Roberts & Marston 2000).

Interval between floods


River Red Gums in EVCs typical of the flood dependent understorey water regime class (Grassy Riverine Forest EVC 106, Riverine Swamp Forest EVC 814 and Sedgy Riverine Forest EVC 816) have a critical interval between flood events ranging from 3-5 years (Fitzsimons et al. 2011). The cumulative effect of repeated dry spells in River Red Gum with flood dependent understorey, for example at more frequent or for longer periods, is unknown. In some circumstances, and on some floodplains, River Red Gums may be largely dependent on water other than flood (surface) water, notably groundwater and/or ponded surface water (Roberts & Marston 2000).

Depth of flooding

Complete immersion, unless brief, is likely to kill River Red Gum seedlings; lower leaves of small saplings die if submerged for long periods. In general, tolerance of flooded conditions increases as seedlings become established, as root system extends and as sapling height increases (Roberts & Marston 2000). Thus two month old seedlings can survive waterlogging for one month with no obvious effect on leaf height and leaf number (Marcar 1993). Seedlings 50-60 cm tall can survive extended flooding of 4-6 months and complete immersion for a few weeks, by shedding leaves (Dexter 1978).

Regarding understorey species, the following inundation depth requirements have been documented:

- *Lobelia concolor* occurs in northern Victoria where flooding is shallow, less than 10cm deep (Ward 1996). However, it would appear to grow in sites which are subject to deeper immersion for short periods (D. Flood, pers. comm. August 2014).
- *Triglochin procerum* typically occurs in depths of 50cm but up to 1.5m. It has been found to tolerate water level increases from 0 to 50cm, from 50-100cm and from 0-100cm (Roberts and Marston 2000).
- *Eleocharis acuta* requires shallow depths, typically 10cm as preferred conditions during its period of active growth. In a glasshouse experiment, it grew 3-4 times better at 0cm depth than under 15cm (Ward 1996; Blanch and Brock 1994; Roberts and Marston 2000). However, this species does tolerate periods of deep inundation (D. Flood, pers. comm. August 2014).



- *Paspalidium jubiflorum* (Warrego Summer Grass) is found where depths are unlikely to exceed 60cm (Roberts and Marston 2000).
- *Carex tereticaulis* tolerates flooding to 10cm in northern Victoria (Roberts and Marston 2000).
- *Phragmites australis* - British studies suggest *Phragmites* tolerates a maximum depth of 0.75 to 1.5 m, and suggests this may be deeper in warmer climates. The plant is tolerant of being overtopped in floods, although prolonged immersion will kill stems. It is also tolerant of fluctuating water levels, and can survive dry conditions. Growth is most vigorous when the water is only a few centimetres deep, and on the banks of the River Murray in South Australia, *Phragmites* occurs where depth is less than 60 cm (Roberts and Marston 2000).
- *Typha domingensis* will grow in areas where water is not too deep (ie < 2 m). On the banks of the River Murray, South Australia, *Typha domingensis* is found mainly in water depths of 20-60 cm (Roberts and Marston 2000).
- *Juncus ingens* in northern Victoria, occurs in depths of up to 1.5 m (Ward 1996).

Native fish (in channel access to floodplain outflows from the forest)

Overview

Describing the hydrological requirements of native fish depends in part on the habitat preference of each species (e.g. channel only, channel and floodplain wetland, off-channel wetland only).


This section focuses on the *requirements of channel specialist native fish*. This mainly includes the large bodied native fish that spawn within channel habitats (such as Murray Cod, Golden Perch and Silver Perch).

The hydrological requirements of native fish that require connectivity with and use of floodplain wetlands have not been included here as (from a fish perspective) the project primarily aims to supply channel fish with productive floodwater inputs that may promote in-channel recruitment. The forest does not have any permanent wetland areas to provide long lasting native fish habitat.

Flow requirements for in-channel recruitment

The flow requirements for recruitment in channel environments, varies among different types of fish (Mallen-Cooper et al. 2013):

- Low-flow channel recruitment (generalists) - Recruitment occurs within channel habitats at low stable flows. Presently only applies to generalist species which, apart from freshwater catfish and olive perchlet, remain common in regulated rivers in the Murray-Darling Basin. The basis of low-flow channel recruitment is that slow-flowing streams with dense plankton enhance survival of fish larvae. The evidence for low-flow channel recruitment was evident in the last drought where the generalist small-bodied species remained abundant in the River Murray. There is also evidence for low-flow recruitment of a range of fish species from the more arid inland river systems (Ebner et al. 2009; Kerezszy et al. 2011).
- Variable flow channel recruitment (channel specialists) - Recruitment occurs when there is variation of within-channel flows. Applies to golden perch, silver perch, and possibly Murray cod and trout cod. In the River Murray variable flow channel recruitment is known to occur for Golden perch and Silver



perch and may be related to increased productivity associated with inundated riverine banks and benches (Mallen-Cooper and Stuart 2003; Zampatti and Leigh 2013). In the Darling River golden perch can spawn during low variable flows (Balcombe et al. 2006; Ebner et al. 2009). These perch species are commonly referred to as flow recruitment specialists (Lake 1967a; Gehrke 1997) as it is generally observed that pulses are needed to generate a spawning response in these species (Mallen-Cooper and Stuart 2003; King et al. 2009). Adult fish develop ova in response to increasing water temperatures (Lake 1967b; Leigh and Zampatti 2011) but can delay spawning, or not spawn at all, if conditions are unsuitable (Leigh and Zampatti 2011) (Baumgartner 2013).

Temperature requirements for in-channel recruitment

In general, the only species that require a rise in flow or a flood to spawn are the flow-cued spawners, golden perch and silver perch. All other species reliably spawn each year in response to rising temperatures in spring and summer (Mallen-Cooper et al. 2013).


This means that to maximise the contribution of floodplain water inputs to in-channel recruitment for species such as golden perch, the floodplain outflows should occur, where possible, after variable channel flows that trigger spawning. For other species, the floodplain outputs should occur in spring/summer after temperature cued spawning has occurred.

Golden Perch

Golden Perch (and Silver Perch) spawn in spring and early summer, in response to rising water temperature and rising flow (Lake 1967, King et al. 2007). Spawning mainly occurs in flowing water habitats in the main channel of rivers and eggs and larvae drift downstream (King et al. 2003, and King et al. 2007). Strong recruitment of these two species is linked to rising flows in spring (Mallen-Cooper and Stuart 2003; Ecological Associates 2010). The River Murray at Torrumbarry Weir is a stronghold for Silver Perch and to a lesser degree Golden Perch where adults and juveniles migrate each year in response to rising flows and water temperatures (Mallen-Cooper 1999).

Golden Perch spawning is associated with high flow events (Reynolds 1983; Hutchinson et al. 2008) and backwater inundation, but recruitment is also evident following a period of zero flow. It exhibits typical flood-induced recruitment, but also opportunistic spawning related to intermittent flooding and dry periods (Balcombe et al. 2006). Golden Perch is regarded as a flood spawner since it tends to spawn and recruit following flow rises, and major spawning occurs when floodplains become inundated (Young et al. 2003; King et al. 2009).

Zampatti and Leigh (2013) found successful spawning and recruitment of Golden Perch occurred during a small but prolonged within-channel increase in discharge in spring/ summer. Most Golden Perch spawned on the ascending limb of the second flow peak of one September to December event, however a few spawned during a small decrease in discharge between two flow peaks. Spawning occurred during a rise and fall in the hydrograph. Hence, some variation in the magnitude of flows rather than a sustained increase in flow may be important to stimulate spawning in Golden Perch. Flows delivered in late spring through summer of >15,000 ML/day when temperatures have reached at least 20°C may promote spawning and recruitment of golden perch and increase population resilience (Zampatti & Leigh 2013).



During spring migration it is possible that movement is directed to specific spawning areas. Evidence for this comes from the movement of fish to distinct reaches within the river. Long distance movements generally occurred between September and December (O'Connor et al. 2005).

Silver Perch

Rogers & Ralph (2011) found the Silver Perch spawning period is from November to January, but is flexible and depends on rising water levels and temperatures. An increase of depth as little as 15cm, in a pond environment can induce spawning (Lake, 1967a). Spawning activity has been found to significantly increase during a flood (Lintermans 2009). Silver Perch migrate upstream to areas behind the peak of a flood, or do so coinciding with a slight rise in water levels and water temperatures (Allen et al. 2003; Lintermans 2007).

Silver Perch is typically regarded as a flood spawner since it tends to spawn and recruit following rises in flow; major spawning occurs when floodplains become inundated (Young et al. 2003).

Murray Cod

Murray Cod spawn around October to December (Humphries 2005; King et al. 2009). A rising spring flow is needed to initiate movement and courtship (Ivor Stuart pers. comm. April 2014). The greatest amount of movement of adult Murray cod has been correlated with peak flows (Koehn & Nicol 1998; Hutchinson et al. 2008).

However, there can also be considerable variation in migration within and between populations; for example, Murray cod can migrate in stable flow and not all Murray cod in a population may migrate. In anabranches Murray cod can have a different pattern again, migrating to spawning areas in autumn or winter rather than spring (Saddler et al. 2008; Mallen-Cooper et al. 2013).

Murray cod and Trout cod are known to spawn over a predictable temporal period in response to increasing temperature irrespective of flow (Rowland 1983; Humphries 2005). Provision of flow increases may not be required as a spawning trigger for these species but could be used to maximize breeding opportunities by inundating more spawning habitat (Baumgartner et al. 2013).


Floodwater contributions to recruitment

According to Mallen-Cooper et al. (2014) recruitment of fish is dependent on the survival of larvae, which have a naturally high mortality. Inundated floodplains are very productive with high densities of phytoplankton and zooplankton that provide food of appropriate size and density for fish larvae. For the fish species that spawn within channel habitats, such as Murray cod, golden perch and silver perch, high survival of larvae may occur as floodwaters recede back into channels. This would be an indirect use of the floodplain by channel specialists and it may occur at a small scale of 10s of kilometres or in large floods over scales of 100s of kilometres. The body of research at present indicates that the channel specialists, which are mostly the large-bodied fish species, can spawn and recruit in absence of floods, especially where there are flowing water habitats without barriers to fish passage, but it is likely that floods significantly enhance recruitment.

Waterbirds

Overview

The literature strongly indicates that a dynamic system is most beneficial to native waterbirds:



- It is a diversity of healthy vegetation types across the floodplain that is likely to increase habitat diversity, food sources and therefore the diversity of waterbird species that use a wetland complex (Ecological Associates 2010).
- Natural or artificial waterbodies that offer an array of water depths and vegetation associations tend to have rich communities of invertebrates, and carry higher numbers of species and individuals of waterbirds (Broome and Jarman 1983).
- The cycle of growth and decay and thus greater availability of nutrients in the water column (Baldwin and Mitchell 2000), resulting from regular inundation and exposure of vegetation along wetland margins, or across the wetland bed, is the basis of a complex food web that provides food to the vertebrates that forage in, on and around the water (Baxter et al. 2005), which in turn support breeding events in many species of fish and waterbirds (Crome 1988, Junk et al. 1989, Scott 1997, Ecological Associates 2010).
- For a complex waterbird community to exist in Gunbower Forest wetlands, a mosaic of shallow gently sloping margins as well as deeper water (>30 cm) and a variety of inundated vegetation types are required (Ecological Associates 2010).
- In Gunbower Forest a drying phase across the floodplain subsequent to inundation is important to waterbird carrying capacity and species diversity (Ecological Associates 2010).


Flood frequency

Waterbirds become sexually mature at the age of one or two years and have a life expectancy ranging from three to four years for ducks and up to eight years for larger birds such as ibis and egrets (Scott, 1997). Therefore waterbirds do not need to breed every year to sustain their populations. However, the provision of optimal breeding conditions within the forest will not necessarily guarantee a bird breeding event. Many factors may prevent the initiation of a breeding event at an artificially watered site, for example a lack of climatic cues and more attractive breeding grounds elsewhere (e.g. Koondrook-Perricoota Forest or Barmah Forest). This occurred in 2008 at Barmah Forest where flooding of breeding areas occurred however the birds did not breed (Leslie, 2008).

Recurrent flooding is considered to be an important factor in building food resources within the forests prior to a managed bird breeding event. A large flood event in the year preceding a managed bird breeding event is highly recommended as this will ensure permanent and semi-permanent wetlands have been refreshed and are productive. Wetlands that are already inundated will provide a source population of wetland vegetation and native fish allowing other areas within the forest to be quickly recolonised. Therefore in planning a managed bird breeding event, it would be desirable to implement a moderate to large watering event in the preceding year (North Central CMA 2009).

Duration

Before waterbirds can breed successfully they need to build up their fat reserves and this can only be achieved if there is sufficient food available. Therefore there is a lag time between the start of a flood and when the birds begin to breed. Warmer temperatures during a spring or early summer flood will result in a faster build up of food resources and therefore the lag time is generally shorter (2 - 3 months). The lag time for an autumn



or winter flood is generally 3 – 6 months, with breeding commencing with the onset of spring (North Central CMA 2009).

Different species have different lag times, all of which are closely related to the type of food each individual species requires and where in the wetland food web the food resource enters the wetland and builds in number. For example ducks will breed relatively quickly requiring 1-2 months of lag time as they graze on algae and invertebrates, which are available at the start of the food web (i.e. when wetland sediments are first inundated) (North Central CMA 2009).

However, egrets require a much longer lag time, up to 7 months, as they are a piscivorous species. Native fish are one of the last food resources to build in number when the wetland is inundated. Therefore the egrets need to wait for the number of fish within the wetland to build so they have enough food resource to last the entire breeding cycle. Experience from flood events at Barmah Forest suggests that an earlier flooding onset will not stimulate birds such as egrets to breed earlier and therefore the key driver is the availability of the food resource (North Central CMA 2009).

To calculate the required duration of a flood event the lag time and the time needed to build nests, lay and incubate eggs and to fledge young (breeding time) is added together. Different species require a different length of time to complete their breeding cycles. Below is a selection of waterbirds common to Benwell Forest and their different breeding timeframes (North Central CMA 2009).

Table 11. Waterbirds typical of Benwell Forest and their different breeding timeframes (North Central CMA 2009)

Waterbird	Lag time (mths)	Breeding time (mths)*	Minimum duration of flooding required (mths)
Ducks	2-3	2-3	4-6
Cormorants	2-3	3-4	5-7
Egrets	7-9	3	10-12

* egg laying, hatching and fledging


Other supporting evidence:

Areas inundated in late winter / early spring must persist for a minimum of 4 (rapid breeders e.g. ducks) to 7 months for the successful breeding of most waterbird species (Scott 1997, Kingsford and Auld 2005). This is because flooding prior to the end of winter will not immediately initiate bird breeding, and this lag needs to be added to the time required for flooding by each species. Consequently, with regard to waterbird breeding, the effective flood duration must be measured from the seasonal increase in water temperatures and concomitant aquatic productivity at the start of spring (Ecological Associates 2010).

Colonial stick-nesting waterbird breeding is dependent on areas of living River Red Gum that are flooded for at least four months (Briggs et al. 1997).

Interval between floods

Crome (1988) found that breeding for a wide variety of waterbird species only followed a rise in water level if the wetland had been completely dried out before hand. This is not true for all waterbird species with Pacific Herons and Yellow-billed Spoonbills favouring sites that have not dried out before reflooding (Briggs et al. 1997).



A drying phase across the floodplain subsequent to inundation is important to waterbird carrying capacity and species diversity. Water bodies that are permanent, or ephemeral systems that lose their drying phase, have been shown to support a: lower density and diversity of birds; decline in invertebrate productivity; increase in abundance of introduced fish; and increase the anaerobic decomposition of organic matter (Crome 1988, Kingston et al. 2004, Gawne and Scholz 2006).

Timing

For most Australian waterbirds, breeding occurs when their food resources are approaching, or are at, a maximum (Kingsford and Norman 2002). The time of year in which flooding occurs is critical for many waterbird species. Floods in winter rarely result in immediate breeding; with many species only initiating breeding in the spring as conditions warm and food resources increase (Loyn et al. 2002; Ecological Associates 2010). The lag between flooding onset and the initiation of breeding relates to such factors as the time required for:


- a large and complex food web to develop, which is capable of supplying abundant food to allow birds to increase fat reserves and develop eggs;
- birds to prepare behaviourally; and
- hormone cycles to be initiated (Ecological Associates 2010).

Waterbirds that feed on animals lower in the food chain (e.g. Ibis feeding on invertebrates) can usually initiate breeding earlier than piscivores (e.g. Darter, Little Black Cormorant, and Intermediate Egrets), which require time for the fish population to develop (Crome 1988). The minimum lag from flood onset to breeding onset in most waterbird species is in the order of 2-3 months (Scott 1997), with breeding of most colonial birds in the Macquarie Marshes positively related to flow and wetland area in the three months before breeding (Kingsford and Auld 2005). This lag time and the causative factors behind it are poorly understood (Ecological Associates 2010).

The most successful waterbird breeding events occur following a flood in late winter, spring or early summer (Scott, 1997), with nesting beginning after birds have had enough time to consume sufficient wetland biota (invertebrates, small fish and aquatic plants) to build up fat reserves (North Central CMA 2009).

To increase the likelihood of waterbirds breeding successfully from an artificial watering event in the Benwell Forest, the flooding should be timed to occur in conjunction with climatic cues. Waterbirds have shown a markedly greater breeding response in these situations than where the flooding occurs independently of such cues (Keith Ward GBCMA, pers. comm. 10/11/2009). Climatic cues include:

- High rainfall in the catchment – will influence available environmental water allocation and potentially influence bird behaviour.
- Natural flood events – taking advantage of natural high rivers and natural flood events (i.e. ‘piggy-backing’ on natural event to extend flood duration) is likely to increase the chance of a successful breeding event.
- Flooding at neighbouring sites – flooding at Guttrum Forest, Campbells Island Forest is likely to impact upon the behaviour of the birds. For example, breeding waterbirds within Gunbower Forest have



been reported to move on a daily basis to the adjacent Koondrook Perricoota area for foraging. Waterbird breeding success is correlated with the foraging area available.

- Irrigation season – colonial waterbirds have been observed to feed in irrigated farm land such as pasture. Success of a breeding event is highly dependant upon a good food resource for the birds. Therefore a large area of irrigated land may contribute to a successful breeding event (North Central CMA 2009).

Depth

Change in depth is an important aspect of waterbird hydrological requirements:

- Many waterbird species will not breed in wetlands with highly controlled water regimes where for example water levels are held at constant levels for extended periods, or alternatively are subject to rapid and/or erratic changes in depth (Briggs et al. 1997);
- inundation and exposure of wetlands needs to occur over seasonal and annual time frames, as long-term rapid and/or erratic changes in water levels within a wetland can result in low numbers of aquatic invertebrates (Briggs et al. 1997), the food of many waterbirds; and
- nearly all colonial nesting waterbirds are vulnerable to sudden drops in water level beneath nesting sites or in foraging areas as discussed above.

In Barmah Forest, a predicted colonial bird breeding event occurred because of the suitable constant low-level flooding through the traditional breeding sites, despite the otherwise drought conditions in the region. Although the higher level transfers were concluded by early December 2002, appropriate ponding levels were maintained by supplying targeted low flows into early February until the colony of predominantly Australian White Ibis and Royal Spoonbill successfully fledged (O'Connor & Ward 2003; Ward 2005). Although targeted water management activities have achieved some good waterbird breeding results over the past decade in Barmah Forest, other breeding attempts have been known to fail due to premature flood subsidence or lack of appropriate flooding (O'Connor & Ward 2003; Webster 2004).

Darters and Cormorants are predominately fish eaters and require open water between 0.6m and 2m deep to obtain food (Ecological Associates 2010).

Wading waterbirds predominantly forage in water up to a maximum depth of approximately 30 cm. Worldwide the greatest diversity and abundance of foraging waterbirds is found in water depths of between 10 and 20 cm (Isola et al. 2000, Taft et al. 2002). Natural or artificial waterbodies that offer an array of water depths and vegetation associations tend to have rich communities of invertebrates, and carry higher numbers of species and individuals of waterbirds (Broome and Jarman 1983). While larger birds will use deeper water in which to forage, they prefer shallow water when food is available (Gawlik 2002), as it is more profitable using less energy to forage (Lovvorn 1994). Piscivores, for example, feed on fish in shallow water in preference to those in deeper water, and the density of prey at which the birds will stop searching increases with increasing depth – being almost twice as high at 28 cm as it is at 10 cm (Gawlik 2002). Maximising the area inundated up to 30 cm in depth in the Benwell Forest will likely increase waterbird species diversity and numbers able to forage in wetlands.